arXiv:1105.3565v2 [math-ph] 19 May 2011

HYPERDIRE
HYPERgeometric functions DIfferential REduction
MATHEMATICA based packages for differential reduction of generalized hypergeometric functions: now with ${}_pF_{p-1}, F_1, F_2, F_3, F_4$

Vladimir V. Bytev[a,b], Mikhail Yu. Kalmykov[a,b], Bernd A. Kniehl[a]

[a] II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, 22761 Hamburg, Germany
[b] Joint Institute for Nuclear Research,
141980 Dubna (Moscow Region), Russia

Abstract

The HYPERDIRE is a project devoted to creation of a set of Mathematica based programs for differential reduction of hypergeometric functions. The current version includes two parts: one, **pfq**, is relevant to manipulation with hypergeometric functions ${}_{p+1}F_p$, and second, **AppellF1F4**, to manipulations with Appell's hypergeometric functions of two variables F_1, F_2, F_3, F_4. As application, a few multiloop Feynman Diagrams of propagator type are considered.

PROGRAM SUMMARY

Title of program: HYPERDIRE

Version: 1.0.0 *Release*: 1.0.0 *Catalogue number*:

Program obtained from `https://sites.google.com/site/loopcalculations/home`:

E-mail: bvv@jinr.ru

Licensing terms: GNU General Public Licence

Computers: all computers running Mathematica

Operating systems: operation systems running Mathematica

Programming language: `Mathematica`

Keywords: Generalized Hypergeometric Functions, Appell Functions, Feynman integrals.

Nature of the problem: Reduction of Hypergeometric Functions: ${}_pF_{p-1}, F_1, F_2, F_3, F_4$ to set of basis functions.

Method of solution: Differential Reduction Reduction

Restriction on the complexity of the problem: none

Typical running time: Depending on the complexity of problem.

LONG WRITE-UP

1 Introduction

Multiple hypergeometric functions [1–4] play an important role in many branches of science. In particular, a large class of Feynman Diagrams are expressed in terms of Horn-type hypergeometric functions [5].

Let us consider a multiple series:

$$H(\vec{\gamma};\vec{\sigma};\vec{x}) = \sum_{m_1,m_2,\cdots,m_r=0}^{\infty} \left(\frac{\Pi_{j=1}^{K} \Gamma\left(\sum_{a=1}^{r} \mu_{ja} m_a + \gamma_j\right) \Gamma^{-1}(\gamma_j)}{\Pi_{k=1}^{L} \Gamma\left(\sum_{b=1}^{r} \nu_{kb} m_b + \sigma_k\right) \Gamma^{-1}(\sigma_k)} \right) x_1^{m_1} \cdots x_r^{m_r} \;, \qquad (1)$$

with $\mu_{ab}, \nu_{ab} \in \mathbb{Q}$, $\gamma_j, \sigma_k \in \mathbb{C}$. The sequences $\vec{\gamma} = (\gamma_1, \cdots, \gamma_K)$ and $\vec{\sigma} = (\sigma_1, \cdots, \sigma_L)$ are called *upper* and *lower* parameters of the hypergeometric function, respectively.

Let $\vec{e_j} = (0, \cdots, 0, 1, 0, \cdots, 0)$ denote the unit vector with unity in its j^{th} entry, and define $\vec{x}^{\vec{m}} = x_1^{m_1} \cdots x_r^{m_r}$ for any integer multi-index $\vec{m} = (m_1, \cdots, m_r)$. Two functions of type (1) with sets of parameters shifted by unity, $H(\vec{\gamma}+\vec{e_c};\vec{\sigma};\vec{x})$ and $H(\vec{\gamma};\vec{\sigma};\vec{x})$, are related by a linear differential operator:

$$H(\vec{\gamma}+\vec{e_c};\vec{\sigma};\vec{x}) \;=\; \frac{1}{\gamma_c} \left(\sum_{a=1}^{r} \mu_{ca} x_a \frac{\partial}{\partial x_a} + \gamma_c \right) H(\vec{\gamma};\vec{\sigma};\vec{x}) \equiv U^{+}_{[\gamma_c \to \gamma_c+1]} H(\vec{\gamma},\vec{\sigma},\vec{x}) \;. \qquad (2)$$

Similar relations also exist for the lower parameters:

$$H(\vec{\gamma};\vec{\sigma}-\vec{e_c};\vec{x}) \;=\; \frac{1}{\sigma_c - 1} \left(\sum_{b=1}^{r} \nu_{cb} x_b \frac{\partial}{\partial x_b} + \sigma_c - 1 \right) H(\vec{\gamma};\vec{\sigma};\vec{x}) \equiv L^{-}_{[\sigma_c \to \sigma_c - 1]} H(\vec{\gamma};\vec{\sigma};\vec{x}) \;. \; (3)$$

The linear differential operators $U^{+}_{\gamma_c \to \gamma_c+1}$, $L^{-}_{\sigma_c \to \sigma_c - 1}$ are called the *step-up* and *step-down* operators for the upper and lower index, respectively. If additional step-down and step-up operators $U^{-}_{\gamma_c}$, $L^{+}_{\sigma_c}$ satisfying

$$U^{-}_{[\gamma_c+1 \to \gamma_c]} U^{+}_{[\gamma_c \to \gamma_c+1]} H(\vec{\gamma},\vec{\sigma},\vec{x}) = L^{+}_{[\sigma_c - 1 \to \sigma_c]} L^{-}_{[\sigma_c \to \sigma_c - 1]} H(\vec{\gamma},\vec{\sigma},\vec{x}) = H(\vec{\gamma},\vec{\sigma},\vec{x})$$

(*i.e.*, the inverses of $U^{+}_{\gamma_c}$, $L^{-}_{\sigma_c}$) are constructed, we can combine these operators to shift the parameters of the hypergeometric function by any integer. This process of applying $U^{\pm}_{\gamma_c}, L^{\pm}_{\sigma_c}$ to shift the parameters by an integer is called **differential reduction** of a hypergeometric function.

In this way, the Horn-type structure provides an opportunity to reduce hypergeometric functions to a set of basis functions with parameters differing from the original values by integer shifts:

$$P_0(\vec{x}) H(\vec{\gamma}+\vec{k};\vec{\sigma}+\vec{l};\vec{x}) = \sum_{m_1,\cdots,m_p=0}^{\sum |k_i| + \sum |l_i|} P_{m_1,\cdots,m_r}(\vec{x}) \left(\frac{\partial}{\partial x_1} \right)^{m_1} \cdots \left(\frac{\partial}{\partial x_r} \right)^{m_r} H(\vec{\gamma};\vec{\sigma};\vec{x}) \;, \quad (4)$$

where $P_0(\vec{x})$ and $P_{m_1,\cdots,m_p}(\vec{x})$ are polynomials with respect to $\vec{\gamma},\vec{\sigma}$ and $\vec{x}$, and $\vec{k},\vec{l}$ are lists of integers.

Algebraic relations between the functions $H(\vec{\gamma},\vec{\sigma};\vec{x})$ with parameters shifted by integers are called **contiguous relations**. The development of systematic techniques for solution of contiguous relations has a long story. It was started by Gauss who described the reduction for the ${}_2F_1$ hypergeometric function in 1823 [1]. Numerous papers have since been published [6–8] on this problem. An algorithmic solution was found by Takayama in Ref. [9], and those methods have been extended in a later series of publications [10, 11] (see also [12]).

Let us recall that any hypergeometric function can be considered to be the solution of a proper system of partial differential equations (PDEs). In particular, for a Horn-type hypergeometric function, the system of PDEs can be derived from the coefficients of the series. In this case, the ratio of two coefficients can be represented as a ratio of two polynomials,

$$\frac{C(\vec{m}+e_j)}{C(\vec{m})} = \frac{P_j(\vec{m})}{Q_j(\vec{m})} \,, \tag{5}$$

so that the Horn-type hypergeometric function satisfies the following system of differential equations:

$$0 = D_j(\vec{\gamma},\vec{\sigma},\vec{x})H(\vec{\gamma},\vec{\sigma},\vec{x}) = \left[Q_j\left(\sum_{k=1}^{r} x_k \frac{\partial}{\partial x_k}\right)\frac{1}{x_j} - P_j\left(\sum_{k=1}^{r} x_k\frac{\partial}{\partial x_k}\right)\right] H(\vec{\gamma},\vec{\sigma},\vec{x}) \,, \tag{6}$$

where $j = 1,\cdots,r$. It was pointed out in several publications [13–15] that (i) the differential reduction algorithm, Eq. (4), can be applied to reduction of Feynman Diagrams to some subset of basis hypergeometric functions with well-known analytical properties [13, 14]; (ii) the system of differential equations, Eq. (6), can be used also for construction of, so-called, ε-expansion of hypergeometric functions around rational values of parameters via direct solution of the system of differential equations [15]. This is another motivation for creating the package for manipulation with parameters of Horn-type hypergeometric functions.

The aim of this paper is to present the *Mathematica* [16] based package **HYPERDIRE** for the differential reduction of the Horn-type hypergeometric function with arbitrary values of parameters to a set of basis functions. The current version consists of two parts: one, **pfq**, for manipulation with hypergeometric function, ${}_{p+1}F_p$, and the second one, **AppellF1F4**, for manipulations with Appell functions, F_1, F_2, F_3, F_4. The algorithm of differential reduction for other functions can be implemented as an additive module.

In contrast to the recent programs written by members of Computational Particles Physics [17–19], the aim of our package is the manipulation with hypergeometric functions without construction of ε-expansion [20, 21].

The preliminary version of **pfq** was presented in [22] and is available in [23]. The latest version is available in [38].

2 Differential reduction algorithm for generalized hypergeometric function ${}_{p+1}F_p$

2.1 General consideration

Let us consider the generalized hypergeometric function, ${}_pF_q(a;b;z)$, defined around $z=0$ by a series

$$
{}_pF_q(\vec{a};\vec{b};z) \equiv {}_pF_q\left(\begin{array}{c} \vec{a} \\ \vec{b} \end{array} \middle| z \right) = \sum_{k=0}^{\infty} \frac{z^k}{k!} \frac{\Pi_{i=1}^{p}(a_i)_k}{\Pi_{j=1}^{q}(b_j)_k} \; , \tag{7}
$$

where $(a)_k$ is a Pochhammer symbol, $(a)_k = \Gamma(a+k)/\Gamma(a)$. The sequences $\vec{a}=(a_1,\cdots,a_p)$ and $\vec{b}=(b_1,\cdots,b_q)$ are called the upper and lower parameters of hypergeometric functions, respectively. In terms of the operator θ:

$$
\theta = z\frac{d}{dz} \; , \tag{8}
$$

the differential equation for the hypergeometric function ${}_pF_q$ can be written as

$$
\left[z\Pi_{i=1}^{p}(\theta+a_i) - \theta\Pi_{i=1}^{q}(\theta+b_i-1) \right] {}_pF_q(\vec{a};\vec{b};z) = 0. \tag{9}
$$

2.2 Differential reduction

The Differential reduction for these functions was analyzed in details in [14]. Here we remind some of the main relations relevant to our program.

The universal differential operators, Eqs. (2),(3), have the following form:

$$
\begin{aligned}
{}_pF_q(a_1+1,\vec{a};\vec{b};z) &= B^{+}_{a_1}{}_pF_q(a_1,\vec{a};\vec{b};z) = \frac{1}{a_1}\left(\theta+a_1\right){}_pF_q(a_1,\vec{a};\vec{b};z) \; , && (10) \\
{}_pF_q(\vec{a};b_1-1,\vec{b};z) &= H^{-}_{b_1}{}_pF_q(\vec{a};b_1,\vec{b};z) = \frac{1}{b_1-1}\left(\theta+b_1-1\right){}_pF_q(\vec{a};b_1,\vec{b};z) \; , && (11)
\end{aligned}
$$

where the operators $B^{+}_{a_1}(H^{-}_{b_1})$ are called the step-up (step-down) operators for the upper (lower) parameters of hypergeometric functions. This type of operators were explicitly constructed for the hypergeometric function ${}_{p+1}F_p$ by Takayama in Ref. [10]. For completeness, we reproduce his result here:

$$
\begin{aligned}
{}_{p+1}F_p(a_i-1,\vec{a};\vec{b};z) &= B^{-}_{a_i}\;{}_{p+1}F_p(a_i,\vec{a};\vec{b};z) \; , \\
{}_{p+1}F_p(\vec{a};b_i+1,\vec{b};z) &= H^{+}_{b_i}\;{}_{p+1}F_p(\vec{a};b_1,\vec{b};z) \; , && (12)
\end{aligned}
$$

where

$$\begin{aligned}
B^-_{a_i} &= -\frac{a_i}{c_i}\left[t_i(\theta) - z\Pi_{j\neq i}(\theta + a_j)\right]\Big|_{a_i\to a_i-1} \ , \\
c_i &= -a_i\Pi^p_{j=1}(b_j - 1 - a_i) \ , \\
t_i(x) &= \frac{x\Pi^p_{j=1}(x + b_j - 1) - c_i}{x + a_i} = \sum_{j=0}^{p} P^{(p)}_{p-j}(\{b_r - 1\})\frac{[x^{j+1} - (-a_i)^{j+1}]}{x + a_i} \\
&= \sum_{j=0}^{p} P^{(p)}_{p-j}(\{b_r - 1\})\sum_{k=0}^{j} x^{j-k}(-a_i)^k \ , \qquad (13)
\end{aligned}$$

and

$$\begin{aligned}
H^+_{a_i} &= \frac{b_i - 1}{d_i}\left[\frac{d}{dz}\Pi_{j\neq i}(\theta + b_j - 1) - s_i(\theta)\right]\Bigg|_{b_i\to b_i+1} \ , \\
d_i &= \Pi^{p+1}_{j=1}(1 + a_j - b_i) \ , \\
s_i(x) &= \frac{\Pi^{p+1}_{j=1}(x + a_j) - d_i}{x + b_i - 1}\sum_{j=0}^{p+1} P^{(p+1)}_{p+1-j}(\{a_r\})\frac{[x^j - (1 - b_i)^j]}{x - (1 - b_i)} \\
&= \sum_{j=0}^{p} P^{(p+1)}_{p-j}(\{a_r\})\sum_{k=0}^{j} x^{j-k}(1 - b_i)^k, \qquad (14)
\end{aligned}$$

where $|_{a\to a+1}$ means substitution of a by $a + 1$, and polynomials $P^{(p)}_j(r_1, \cdots, r_p)$ are defined as

$$\prod_{k=1}^{p}(z + r_k) = \sum_{j=0}^{p} P^{(p)}_{p-j}(r_1, \cdots, r_p)z^j \equiv \sum_{j=0}^{p} P^{(p)}_{p-j}(\vec{r})z^j \equiv \sum_{j=0}^{p} P^{(p)}_{j}(\vec{r})z^{p-j} \ . \qquad (15)$$

$P^{(p)}_s(\vec{r})$ is a polynomial of an order of s with respect to variables r, and

$$P^{(p)}_0(\vec{r}) = 1 \ , \quad P^{(p)}_j(\vec{r}) = \sum_{i_1,\cdots,i_r=1}^{p}\prod_{i_1<\cdots<i_j} r_{i_1}\cdots r_{i_j} \ , \quad j = 1, \cdots, p \ .$$

For example, $P^{(p)}_1(\vec{r}) = \sum_{j=1}^{p} r_j$ and $P^{(p)}_p(\vec{r}) = \prod_{j=1}^{p} r_j$. Keeping in mind that

$$\prod_{i=1}^{p}(z + r_i)\prod_{j=p+1}^{p+k}(z + r_j) = \prod_{l=1}^{p+k}(z + r_l) \ ,$$

we get that these polynomials satisfy the following relations:

$$P^{(p+k)}_{p+k-j}(r_1, \cdots, r_p, q_1, \cdots, q_k) = \sum_{n=0}^{k} P^{(p)}_{p+1-j-n}(r_1, \cdots, r_p)P^{(k)}_n(q_1, \cdots, q_k) \ , \qquad (16)$$

where $j = 1, \cdots, p-k$. In particular,

$$\begin{aligned}
P^{(p+1)}_{p+1-j}(\vec{r}, f) &= P^{(p)}_{p+1-j}(\vec{r}) + f P^{(p)}_{p-j}(\vec{r}) \;, \quad j = 1, \cdots, p. \\
P^{(p+1)}_{p+1-j}(\vec{r}_{p-1}, q_1, q_2) &= \sum_{k=0}^{2} P^{(p-1)}_{p+1-j-k}(\vec{r}) P^{(2)}_{k}(\vec{q}) \;, \quad j = 1, \cdots, p-1. \\
P^{(p+1)}_{p+1-j}(\vec{r}_{p-2}, q_1, q_2, q_3) &= \sum_{k=0}^{3} P^{(p-2)}_{p+1-j-k}(\vec{r}) P^{(3)}_{k}(\vec{q}) \;, \quad j = 1, \cdots, p-2.
\end{aligned}$$

The differential reduction has the form of a product of several differential step-up/step-down operators, $H^{\pm}_{b_k}$ and $B^{\pm}_{a_k}$, respectively:

$$F(\vec{a} + \vec{m}; \vec{b} + \vec{n}; z) = \left(H^{\pm}_{\{a\}} \right)^{\sum_i m_i} \left(B^{\pm}_{\{b\}} \right)^{\sum_j n_j} F(\vec{a}; \vec{b}; z) \;, \tag{17}$$

so that maximal power of θ in this expression is equal to $r \equiv \sum_i m_i + \sum_j n_j$. Since the hypergeometric function ${}_{p+1}F_p(\vec{a}; \vec{b}; z)$ satisfies the differential equation of an order of $p+1$ (see Eq. (6)):

$$\begin{aligned}
&(1-z)\theta^{p+1} {}_{p+1}F_p(\vec{a}; \vec{b}; z) \\
&= \left\{ \sum_{r=1}^{p} \left[z P^{(p+1)}_{p+1-r}(\{a_j\}) - P^{(p)}_{p+1-r}(\{b_j - 1\}) \right] \theta^r + z \Pi_{k=1}^{p+1} a_k \right\} {}_{p+1}F_p(\vec{a}; \vec{b}; z) \;,
\end{aligned} \tag{18}$$

it is possible to express all terms containing higher power of the operator θ^k, where $k \geq p+1$, in terms of rational functions of parameters and argument z times by θ^j, where $j \leq p$.

In this way, any function ${}_{p+1}F_p(\vec{a}+\vec{m}; \vec{b}+\vec{k}; z)$ is expressible in terms of the basic function and its first p-derivative:

$$\begin{aligned}
&{}_{p+1}F_p(\vec{a} + \vec{m}; \vec{b} + \vec{k}; z) = \\
&\frac{1}{S(a_i, b_j, z)} \left\{ R_1(a_i, b_j, z) + R_2(a_i, b_j, z)\theta + \cdots + R_{p+1}(a_i, b_j, z)\theta^p \right\} {}_{p+1}F_p(\vec{a}; \vec{b}; z) \;,
\end{aligned} \tag{19}$$

where m, k is the set of integer numbers, and S and R_i are polynomials in the parameters $\{a_i\}, \{b_j\}$ and z.

From Eq. (13) follows that if one of the upper parameters a_j is equal to unity, then the application of the step-down operator $B^{-}_{a_j}$ to the hypergeometric function ${}_{p+1}F_p$ will produce unity, $B^{-}_{1} \, {}_{p+1}F_p(1, \vec{a}; \vec{b}; z) \equiv 1$. Taking into account the explicit form of the step-down operator B^{-}_{1},

$$B^{-}_{1} = \frac{1}{\Pi_{k=1}^{p}(b_k - 1)} \left[\Pi_{j=1}^{p}(b_j - 1) + \sum_{j=1}^{p} P^{(p)}_{p-j}(\{b_k - 1\})\theta^j - z \Pi_{j=1}^{p} (\theta + a_j) \right] \;, \tag{20}$$

we get the differential identity

$$\left\{ \Pi_{j=1}^{p}(b_j-1) - z\Pi_{j=1}^{p} a_j + (1-z)\theta^p \right\}{}_{p+1}F_p(1,\vec{a};\vec{b};z)$$
$$+\left\{ \sum_{j=1}^{p-1} \left[P_{p-j}^{(p)}(\{b_k-1\}) - zP_{p-j}^{(p)}(\{a_k\}) \right] \theta^j \right\}{}_{p+1}F_p(1,\vec{a};\vec{b};z) = \Pi_{k=1}^{p}(b_k-1) \ . \quad (21)$$

The case when two or more upper parameters are equal to unity, $a_1 = a_2 = 1$, does not generate any new identities.

3 Differential reduction for the Appell hypergeometric functions

3.1 Appell hypergeometric functions: system of differential equations

Let us consider the system of linear differential equations of the second order for the functions $\omega(\vec{z})$:

$$\theta_{11}\omega(\vec{z}) = \left\{ P_0(\vec{z})\theta_{12} + P_1(\vec{z})\theta_1 + P_2(\vec{z})\theta_2 + P_3(\vec{z}) \right\}\omega(\vec{z}) \ , \quad (22)$$

$$\theta_{22}\omega(\vec{z}) = \left\{ R_0(\vec{z})\theta_{12} + R_1(\vec{z})\theta_1 + R_2(\vec{z})\theta_2 + R_3(\vec{z}) \right\}\omega(\vec{z}) \ , \quad (23)$$

where $\vec{z} = (z_1, z_2)$ and z_1, z_2 are variables, $\{P_j, R_j\}$ are rational functions, $\theta_j = z_j\partial_{z_j}$ for $j = 1,2$ and $\theta_{i_1\cdots i_k} = \theta_{i_i}\cdots\theta_{i_k}$. The exploring of θ_j instead of the standard ∂_j is explained by our applications. Taking the derivative of Eq.(22) with respect to θ_2, using the well-known property $\partial_2\partial_{11}\omega(\vec{z}) = \partial_1\partial_{12}\omega(\vec{z})$ and applying Eq. (23), we rewrite the first equation as follows:

$$[\theta_1 - P_0\theta_2]\,\theta_{12}\omega(\vec{z})$$
$$= \left\{ [\theta_2 P_0 + P_1 + P_2 R_0]\,\theta_{12} + [P_2 R_1 + \theta_2 P_1]\,\theta_1 + [P_2 R_2 + \theta_2 P_2 + P_3]\,\theta_2 + P_2 R_3 + \theta_2 P_3 \right\}\omega(\vec{z}) \ , \quad (24)$$

Repeating a similar operation for Eq. (23) we get

$$[-R_0\theta_1 + \theta_2]\,\theta_{12}\omega(\vec{z})$$
$$= \left\{ [\theta_1 R_0 + R_2 + R_1 P_0]\,\theta_{12} + [P_1 R_1 + \theta_1 R_1 + R_3]\,\theta_1 + [P_2 R_1 + \theta_1 R_2]\,\theta_2 + R_1 P_3 + \theta_1 R_3 \right\}\omega(\vec{z}) \ , \quad (25)$$

It is well known [2] that under condition

$$1-P_0R_0\neq 0\ , \tag{26}$$

there are 4 independent solutions of system (22),(23). In this case, Eqs. (24),(25) can be solved so that

$$\begin{aligned}
(1-P_0R_0)\theta_{112}\omega(\vec{z}) \;=\;& \left\{ [P_0\left(\theta_1R_0+R_2+R_1P_0\right)+\theta_2P_0+P_1+P_2R_0]\,\theta_{12}^2 \right. \\
&+[P_2R_1+\theta_2P_1+P_0\left(P_1R_1+\theta_1R_1+R_3\right)]\,\theta_1 \\
&+[P_2R_2+\theta_2P_2+P_3+P_0\left(P_2R_1+\theta_1R_2\right)]\,\theta_2 \\
&\left. +P_2R_3+\theta_2P_3+P_0\left(R_1P_3+\theta_1R_3\right) \right\}\omega(\vec{z})\ , \qquad (27) \\
(1-P_0R_0)\theta_{122}\omega(\vec{z}) \;=\;& \left\{ [\theta_1R_0+R_2+R_1P_0+R_0\left(\theta_2P_0+P_1+P_2R_0\right)]\,\theta_{12}^2 \right. \\
&+[R_0\left(P_2R_1+\theta_2P_1\right)+P_1R_1+\theta_1R_1+R_3]\,\theta_1 \\
&+[P_2R_1+\theta_1R_2+R_0\left(P_2R_2+\theta_2P_2+P_3\right)]\,\theta_2 \\
&\left. +R_0\left(P_2R_3+\theta_2P_3\right)+R_1P_3+\theta_1R_3 \right\}\omega(\vec{z})\ . \qquad (28)
\end{aligned}$$

The condition of complete integrability is defined from the relation $\partial_1\left(\partial_{122}\omega(\vec{z})\right)=\partial_2\left(\partial_{112}\omega(\vec{z})\right)$, or in terms of the quantities (27) and (28) it has the following form:

$$\begin{aligned}
&\theta_2\left(\frac{1}{1-P_0R_0}\,[\text{r.h.s.Eq. (27)}]\right)-\theta_1\left(\frac{1}{1-P_0R_0}\,[\text{r.h.s.Eq. (28)}]\right) \\
&=\frac{1}{1-P_0R_0}\,[\text{r.h.s.Eq. (27)}]-\frac{1}{1-P_0R_0}\,[\text{r.h.s.Eq. (28)}] \\
&\quad+\theta_2\left(\theta_{12}\omega(\vec{z})\right)-\theta_1\left(\theta_{12}\omega(\vec{z})\right)\ ,
\end{aligned} \tag{29}$$

where we have used the following relation $\theta_{jj}=z_j^2\partial_j^2+z_j\partial_j$. If the condition (29) is valid, Eqs. (22) and (23) can be reduced to the Pfaff system of four differential equations

$$d\vec{f}=R\vec{f}\ , \tag{30}$$

where $\vec{f}=(\omega(\vec{z}),\theta_1\omega(\vec{z}),\theta_2\omega(\vec{z}),\theta_{12}^2\omega(\vec{z}))$.

In the case

$$1-P_0R_0=0\ , \tag{31}$$

$\theta_{12}\omega(\vec{z})$ are expressible in terms of three other elements: $\omega(\vec{z}),\theta_1\omega(\vec{z}),\theta_2\omega(\vec{z})$. In particular,

Table 1:

	F_1	F_2	F_3	F_4
P_0	- 1	$\frac{z_1}{1-z_1}$	$-\frac{1}{(1-z_1)}$	$\frac{2z_1}{(1-z_1-z_2)}$
R_0	- 1	$\frac{z_2}{1-z_2}$	$-\frac{1}{(1-z_2)}$	$\frac{2z_2}{(1-z_1-z_2)}$
P_1	$\frac{(a+b_1)z_1-(c-1)}{(1-z_1)}$	$\frac{(a+b_1)z_1-(c_1-1)}{(1-z_1)}$	$\frac{(a_1+b_1)z_1-(c-1)}{(1-z_1)}$	$\frac{(a+b)z_1-(c_1-1)(1-z_2)}{(1-z_1-z_2)}$
R_1	$\frac{b_2z_2}{(1-z_2)}$	$\frac{b_2z_2}{(1-z_2)}$	0	$\frac{(a+b+1-c_1)z_2}{(1-z_1-z_2)}$
P_2	$\frac{b_1z_1}{(1-z_1)}$	$\frac{b_1z_1}{(1-z_1)}$	0	$\frac{(a+b+1-c_2)z_1}{(1-z_1-z_2)}$
R_2	$\frac{(a+b_2)z_2-(c-1)}{(1-z_2)}$	$\frac{(a+b_2)z_2-(c_2-1)}{(1-z_2)}$	$\frac{(a_2+b_2)z_2-(c-1)}{(1-z_2)}$	$\frac{(a+b)z_2-(c_2-1)(1-z_1)}{(1-z_1-z_2)}$
P_3	$\frac{ab_1z_1}{(1-z_1)}$	$\frac{ab_1z_1}{(1-z_1)}$	$\frac{a_1b_1z_1}{(1-z_1)}$	$\frac{abz_1}{(1-z_1-z_2)}$
R_3	$\frac{ab_2z_2}{(1-z_2)}$	$\frac{ab_2z_2}{(1-z_2)}$	$\frac{a_2b_2z_2}{(1-z_2)}$	$\frac{abz_2}{(1-z_1-z_2)}$

in terms of the notation of Eqs.(24), (25),

$$\begin{aligned} & \left[\theta_2 P_0 + P_1 + P_2 R_0 + P_0\left(\theta_1 R_0 + R_2 + R_1 P_0\right)\right]\theta_{12}\omega(\vec{z}) \\ = & \left\{ - \left[P_0\left(\theta_1 R_1 + R_3 + R_1 P_1\right) + P_2 R_1 + \theta_2 P_1\right]\theta_1 \right. \\ & - \left[P_0\left(\theta_1 R_2 + R_1 P_2\right) + P_2 R_2 + \theta_2 P_2 + P_3\right]\theta_2 \\ & \left. - \left[P_0\left(\theta_1 R_3 + R_1 P_3\right) + P_2 R_3 + \theta_2 P_3\right] \right\} \omega(\vec{z}) \, . \end{aligned} \tag{32}$$

In this case, the integrability conditions are valid and Eqs. (22) and (23) can be reduced to the Pfaff system (30) of three differential equations $\vec{f} = (\omega(\vec{z}), \theta_1\omega(\vec{z}), \theta_2\omega(\vec{z}))$, and the system has 3 solutions.

For Appell's hypergeometric functions F_1, F_2, F_3 and F_4 the values of coefficients of Eqs. (22), (23) are collected in Table 1.

3.2 Appell hypergeometric function: F_1

3.2.1 General consideration

Let us consider the Appell hypergeometric functions F_1 defined around $x = y = 0$ as

$$\omega \equiv F_1(a, b_1, b_2, c; x, y) = \sum_{m=0}^{\infty}\sum_{n=0}^{\infty} \frac{(a)_{m+n}(b_1)_m(b_2)_n}{(c)_{m+n}} \frac{x^m}{m!}\frac{y^n}{n!} \, . \tag{33}$$

In this case, Eqs. (22) and (23) have the following form:

$$\theta_{xx}\omega = \left[\frac{(a+b_1)x-(c-1)}{1-x} - b_2\frac{y}{x-y}\right]\theta_x\omega + \frac{b_1x(1-y)}{(1-x)(x-y)}\theta_y\omega + \frac{x}{1-x}ab_1\omega \, , \tag{34}$$

$$\theta_{yy}\omega = \left[\frac{(a+b_2)y-(c-1)}{1-y} + b_1\frac{x}{x-y}\right]\theta_y\omega - \frac{b_2y(1-x)}{(1-y)(x-y)}\theta_x\omega + \frac{y}{1-y}ab_2\omega \, . \tag{35}$$

The Eq. (31) is fulfilled and Eq. (32) has the following form:

$$(x-y)\frac{\partial^2\omega}{\partial x\partial y} - b_2\frac{\partial\omega}{\partial x} + b_1\frac{\partial\omega}{\partial y} = 0 \; , \tag{36}$$

or in terms of the operators θ_x, θ_y

$$\theta_{xy}\omega = \frac{b_2 y}{x-y}\theta_x\omega - \frac{b_1 x}{x-y}\theta_y\omega \; . \tag{37}$$

3.2.2 Differential Reduction of F_1

The direct differential expressions follow from Eqs. (2) and (3)

$$\begin{aligned} aF_1(a+\mathbf{1},b_1,b_2,c;x,y) &= (\theta_x{+}\theta_y{+}a)F_1(a,b_1,b_2,c;x,y) \; , & (38)\\ b_1F_1(a,b_1+\mathbf{1},b_2,c;x,y) &= (\theta_x{+}b_1)F_1(a,b_1,b_2,c;x,y) \; , & (39)\\ (c-1)F_1(a,b_1,b_2,c-\mathbf{1};x,y) &= (\theta_x{+}\theta_y{+}c{-}1)F_1(a,b_1,b_2,c;x,y) \; . & (40) \end{aligned}$$

The inverse differential relations were considered in [2, 7]:

$$\begin{aligned} &(c{-}a)F_1(a-\mathbf{1},b_1,b_2,c;x,y) = \\ &\left[c{-}a{-}b_1x{-}b_2y{+}(1{-}x)\theta_x{+}(1{-}y)\theta_y\right]F_1(a,b_1,b_2,c;x,y) \; , && (41)\\ &(c{-}b_1{-}b_2)F_1(a,b_1-\mathbf{1},b_2,c;x,y) = \\ &\left[c{-}b_1{-}b_2{-}ax{+}(1{-}x)\theta_x{-}x\left(1{-}\frac{1}{y}\right)\theta_y\right]F_1(a,b_1,b_2,c;x,y) \; , && (42)\\ &(c{-}a)(c{-}b_1{-}b_2)F_1(a,b_1,b_2,c+\mathbf{1};x,y) = \\ &c\left[(c{-}a{-}b_1{-}b_2){-}\left(1{-}\frac{1}{x}\right)\theta_x{-}\left(1{-}\frac{1}{y}\right)\theta_y\right]F_1(a,b_1,b_2,c;x,y) \; . && (43) \end{aligned}$$

The differential reduction for the parameter "b_2" follows from Eqs. (39) and (42) and the symmetry property of the function F_1: $F_1(a,b_1,b_2,c;x,y) = F_1(a,b_2,b_1,c;y,x)$, so that

$$b_1 \Leftrightarrow b_2 \; , \quad x \Leftrightarrow y \; .$$

3.3 Appell hypergeometric function: F_2

3.3.1 General consideration

Let us consider the Appell hypergeometric functions F_2 defined around $x = y = 0$ as

$$\omega \equiv F_2(a,b_1,b_2,c_1,c_2;x,y) = \sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\frac{(a)_{m+n}(b_1)_m(b_2)_n}{(c_1)_m(c_2)_n}\frac{x^m}{m!}\frac{y^n}{n!} \; , \tag{44}$$

In this case Eqs. (22) and (23) have the following form:

$$\begin{aligned} (1{-}x)\theta_{xx}\omega &= x\theta_{xy}\omega + \left[(a{+}b_1)x{-}(c_1{-}1)\right]\theta_x\omega{+}b_1x\theta_y\omega{+}ab_1x\omega \; , & (45)\\ (1{-}y)\theta_{yy}\omega &= y\theta_{xy}\omega + \left[(a{+}b_2)y{-}(c_2{-}1)\right]\theta_y\omega{+}b_2y\theta_x\omega{+}ab_2y\omega \; . & (46) \end{aligned}$$

The condition, Eq. (26), is fulfilled and Eqs. (27) and (28) have the following form:

$$\begin{aligned}
(1-x-y)\theta_{xxy}\omega &= \left[(a+b_1+1-c_2)\,x-(c_1-1)(1-y)+\frac{b_2xy}{1-x}\right]\theta_{xy}\omega \\
& +\frac{b_2xy}{1-x}\left[a+b_1+1-c_1\right]\theta_x\omega \\
& +\left[(a+1-c_2)b_1x+\frac{b_1b_2xy}{1-x}\right]\theta_y\omega \\
& +\frac{ab_1b_2xy}{1-x}\omega \;, \qquad (47) \\
(1-x-y)\theta_{yyx}\omega &= \left[(a+b_2+1-c_1)\,y-(c_2-1)(1-x)+\frac{b_1xy}{1-y}\right]\theta_{xy}\omega \\
& +\frac{b_1xy}{1-y}\left[a+b_2+1-c_2\right]\theta_y\omega \\
& +\left[(a+1-c_1)b_2y+\frac{b_1b_2xy}{1-y}\right]\theta_x\omega \\
& +\frac{ab_1b_2xy}{1-y}\omega \;. \qquad (48)
\end{aligned}$$

3.3.2 Differential Reduction of F_2

The direct differential expressions follow from Eqs. (2) and (3)

$$\begin{aligned}
aF_2(a+\mathbf{1},b_1,b_2,c_1,c_2;x,y) &= (a+\theta_x+\theta_y)F_2(a,b_1,b_2,c_1,c_2;x,y)\;, \qquad (49) \\
b_1F_2(a,b_1+\mathbf{1},b_2,c_1,c_2;x,y) &= (b_1+\theta_x)F_2(a,b_1,b_2,c_1,c_2;x,y)\;, \qquad (50) \\
(c_1-1)F_2(a,b_1,b_2,c_1-\mathbf{1},c_2;x,y) &= (c_1-1+\theta_x)F_2(a,b_1,b_2,c_1,c_2;x,y)\;, \qquad (51)
\end{aligned}$$

The inverse differential relations were considered in [7]:

$$\begin{aligned}
& F_2(a-\mathbf{1},b_1,b_2,c_1,c_2;x,y) = \Bigg\{1-\frac{xb_1-(1-x)\theta_x}{c_1-a}-\frac{yb_2-(1-y)\theta_y}{c_2-a} \\
& +\frac{1}{c_1+c_2-a-1}\left[\frac{1}{c_1-a}+\frac{1}{c_2-a}\right]\left[(1-x-y)\theta_{xy}-b_1x\theta_y-b_2y\theta_x\right]\Bigg\} \\
& \times F_2(a,b_1,b_2,c_1,c_2;x,y)\;, \qquad (52) \\
& F_2(a,b_1-\mathbf{1},b_2,c_1,c_2;x,y) = \left\{1+\frac{(1-x)\theta_x-x(a+\theta_y)}{c_1-b_1}\right\}F_2(a,b_1,b_2,c_1,c_2;x,y)\;, \qquad (53) \\
& F_2(a,b_1,b_2,c_1+\mathbf{1},c_2;x,y) = \frac{c_1}{(c_1-a)(c_1-b_1)}\Bigg\{c_1-a-b_1-\left(1-\frac{1}{x}\right)\theta_x \\
& -\frac{1}{x(c_1+c_2-a-1)}\left[xb_1\theta_y+yb_2\theta_x-(1-x-y)\theta_{xy}\right]\Bigg\}F_2(a,b_1,b_2,c_1,c_2;x,y)\;. \qquad (54)
\end{aligned}$$

The differential reduction for the parameters "b_2" and "c_2" follows from Eqs. (50),(53) and Eqs. (51),(54) and symmetry property of function F_2: $F_2(a,b_1,b_2,c;x,y) = F_2(a,b_2,b_1,c;y,x)$:

$$b_1 \Leftrightarrow b_2 \; , \quad c_1 \Leftrightarrow c_2 \; , \quad x \Leftrightarrow y \; .$$

3.4 Appell hypergeometric function: F_3

3.4.1 General consideration

Let us consider the Appell hypergeometric functions F_3 defined around $x = y = 0$ as

$$\omega \equiv F_3(a_1,a_2,b_1,b_2,c;x,y) = \sum_{m=0}^{\infty}\sum_{n=0}^{\infty} \frac{(a_1)_m (a_2)_n (b_1)_m (b_2)_n}{(c)_{m+n}} \frac{x^m}{m!}\frac{y^n}{n!} \; , \tag{55}$$

In this case Eqs. (22) and (23) have the following form:

$$\begin{aligned} (1-x)\theta_{xx}\omega &= -\theta_{xy}\omega + \left[(a_1+b_1)x-(c-1)\right]\theta_x\omega + x a_1 b_1 \omega \; , & (56) \\ (1-y)\theta_{yy}\omega &= -\theta_{xy}\omega + \left[(a_2+b_2)y-(c-1)\right]\theta_y\omega + y a_2 b_2 \omega \; . & (57) \end{aligned}$$

The condition, Eq. (26), is fulfilled and Eqs. (27) and(28) have the following form:

$$\begin{aligned} (xy-x-y)\theta_{xxy}\omega &= \left[(1-y)(a_1+b_1)x - y(a_2+b_2+1-c)\right]\theta_{xy}\omega \\ &\quad + (1-y)x a_1 b_1 \theta_y\omega - y a_2 b_2 \theta_x \omega \; , & (58) \\ (xy-x-y)\theta_{xyy}\omega &= \left[(1-x)(a_2+b_2)y - x(a_1+b_1+1-c)\right]\theta_{xy}\omega \\ &\quad + (1-x)y a_2 b_2 \theta_x\omega - x a_1 b_1 \theta_y \omega \; . & (59) \end{aligned}$$

3.4.2 Differential Reduction of F_3

The direct differential expressions follow from Eqs. (2) and (3)

$$\begin{aligned} a_1 F_3(a_1+\mathbf{1},a_2,b_1,b_2,c;x,y) &= (a_1+\theta_x)F_3(a_1,a_2,b_1,b_2,c;x,y) \; , & (60) \\ b_1 F_3(a_1,a_2,b_1+\mathbf{1},b_2,c;x,y) &= (b_1+\theta_x)F_3(a_1,a_2,b_1,b_2,c;x,y) \; , & (61) \\ (c-1)F_3(a_1,a_2,,b_1,b_2,c-\mathbf{1};x,y) &= (c-1+\theta_x+\theta_y)F_3(a,b_1,b_2,c_1,c_2;x,y) \; . & (62) \end{aligned}$$

The inverse differential relations were considered in [8]:

$$F_3(a_1-\mathbf{1},a_2,b_1,b_2,c;x,y) = 1+\frac{1}{(c-a_1-a_2)(c-b_2-a_1)}$$

$$\left\{(c-b_2-a_1-a_2)\left[(1-x)\theta_x-xb_1\right]+b_1x\left(1-\frac{1}{y}\right)\theta_y-\left(1-x+\frac{x}{y}\right)\theta_{xy}\right\}$$

$$\times F_3(a,b_1,b_2,c_1,c_2;x,y)\ , \tag{63}$$

$$F_3(a_1,a_2,b_1-\mathbf{1},b_2,c;x,y) = 1+\frac{1}{(c-b_1-b_2)(c-a_2-b_1)}$$

$$\left\{(c-a_2-b_1-b_2)\left[(1-x)\theta_x-xa_1\right]+a_1x\left(1-\frac{1}{y}\right)\theta_y-\left(1-x+\frac{x}{y}\right)\theta_{xy}\right\}$$

$$\times F_3(a,b_1,b_2,c_1,c_2;x,y)\ , \tag{64}$$

$$\Delta F_3(a_1,a_2,b_1,b_2,c+\mathbf{1};x,y) =$$

$$c\left\{A-D_1\left(1-\frac{1}{x}\right)\theta_x-D_2\left(1-\frac{1}{y}\right)\theta_y+B\left(1-\frac{1}{x}-\frac{1}{y}\right)\theta_{xy}\right\}$$

$$\times F_3(a,b_1,b_2,c_1,c_2;x,y)\ , \tag{65}$$

where in the last expression,

$$\begin{aligned}
&\delta_1 = c-a_1-b_1\ , \quad \delta_2 = c-a_2-b_2\ , \quad F = c-a_1-a_2-b_1-b_2\ , \\
&A = \delta_1\delta_2F+a_1b_1\delta_1+a_2b_2\delta_2\ , \\
&D_1 = \delta_2F+a_1b_1-a_2b_2\ , \quad D_2 = \delta_1F+a_2b_2-a_1b_1\ , \quad B = \delta_1+\delta_2\ , \\
&\Delta = (c-b_1-b_2)(c-a_1-a_2)(c-a_2-b_1)(c-a_1-b_2)\ .
\end{aligned} \tag{66}$$

The differential reduction for the parameters "a_2" and "b_2" follows from Eqs. (60), (63) and Eqs. (61), (64), respectively, and the symmetry property of the function F_3 :
$F_3(a_1,a_2,b_1,b_2,c;x,y) = F_3(a_2,a_1,b_2,b_1,c;y,x)$:

$$a_1 \Leftrightarrow a_2\ , \quad b_1 \Leftrightarrow b_2\ , \quad x \Leftrightarrow y\ .$$

3.5 Appell hypergeometric function: F_4

3.5.1 General consideration

Let us consider the Appell hypergeometric functions F_4 defined around $x=y=0$ as

$$\omega \equiv F_4(a,b,c_1,c_2;x,y) = \sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\frac{(a)_{m+n}(b)_{m+n}}{(c_1)_m(c_2)_n}\frac{x^m}{m!}\frac{y^n}{n!}\ , \tag{67}$$

The condition, Eq. (26), is fulfilled and Eqs. (27),(28) have the following form:

$$(1-x-y)\theta_{xx}\omega = 2x\theta_{xy}\omega+\left[(a+b)x-(c_1-1)(1-y)\right]\theta_x\omega+\left[a+b+1-c_2\right]x\theta_y\omega+abx\omega\ , \tag{68}$$

$$(1-x-y)\theta_{yy}\omega = 2y\theta_{xy}\omega+\left[(a+b)y-(c_2-1)(1-x)\right]\theta_y\omega+\left[a+b+1-c_1\right]y\theta_x\omega+aby\omega\ . \tag{69}$$

The condition, Eq. (26), is fulfilled. However, instead of Eqs. (27),(28), which are very long in this case, we present the results in the following form:

$$
\begin{aligned}
&\left[(1-x-y)^2-4xy\right]\theta_{xxy}\omega = \\
&y\left[1-x-y+2x(a+b+1-c_1)\right]\theta_{xx}\omega + x\left[2x+(1-x-y)(a+b+1-c_2)\right]\theta_{yy}\omega + \\
&\left[2(a+b+1-c_2)(1-x)x - (c_1-1)(1-x-y)^2 - x(1-x-y)(a+b+c_1-1)\right]\theta_{xy}\omega \\
&+y\left[(c_1-1)(1-x-y)+2abx\right]\theta_{x}\omega + x\left[2x(c_2-1)+(1-x-y)ab\right]\theta_{y}\omega \; ,
\end{aligned}
\tag{70}
$$

$$
\begin{aligned}
&\left[(1-x-y)^2-4xy\right]\theta_{yyx}\omega = \\
&x\left[1-x-y+2y(a+b+1-c_2)\right]\theta_{yy}\omega + y\left[2y+(1-x-y)(a+b+1-c_1)\right]\theta_{xx}\omega + \\
&\left[2(a+b+1-c_1)(1-y)y - (c_2-1)(1-x-y)^2 - y(1-x-y)(a+b+c_2-1)\right]\theta_{xy}\omega \\
&+x\left[(c_2-1)(1-x-y)+2aby\right]\theta_{y}\omega + y\left[2y(c_1-1)+(1-x-y)ab\right]\theta_{x}\omega \; ,
\end{aligned}
\tag{71}
$$

where the values of $\theta_{xx}\omega$ and $\theta_{yy}\omega$ are taken from Eqs. (68) and (69).

3.5.2 Differential Reduction of F_4

The direct differential expressions follow from Eqs. (2) and (3)

$$
aF_4(a+\mathbf{1},b,c_1,c_2;x,y) = (a+\theta_x+\theta_y)F_4(a,b,c_1,c_2;x,y) \; , \tag{72}
$$

$$
(c_1-1)F_4(a,b,c_1-\mathbf{1},c_2;x,y) = (c_1-1+\theta_x)F_4(a,b,c_1,c_2;x,y) \; , \tag{73}
$$

The inverse differential relations were considered in [7]:

$$
\begin{aligned}
& F_4(a-\mathbf{1},b,c_1,c_2;x,y) = \Biggl\{ 1 \\
& -\frac{x}{(c_1-a)}\left[\left(1-\frac{1}{x}\right)\theta_x+\theta_y+b\right]-\frac{y}{(c_2-a)}\left[\left(1-\frac{1}{y}\right)\theta_y+\theta_x+b\right] \\
& +\frac{1}{(1-x-y)(c_1+c_2-a-1)}\left[\frac{1}{c_1-a}+\frac{1}{c_2-a}\right]\Biggl(\\
& \qquad [(1-x-y)^2-4xy]\theta_{xy} \\
& \qquad -y[2x(a+b+1-c_1)+(1-x-y)(b+1-c_1)]\theta_x \\
& \qquad -x[2y(a+b+1-c_2)+(1-x-y)(b+1-c_2)]\theta_y \\
& \qquad -2abxy \qquad\qquad\qquad \Biggr)\Biggr\} F_4(a,b,c_1,c_2;x,y)\ ,
\end{aligned}
\tag{74}
$$

$$
\begin{aligned}
& x(c_1-a)(c_1-b)F_4(a,b,c_1+\mathbf{1},c_2;x,y) = \\
& c_1\Biggl\{ x(c_1-a-b)+(1-x-y)\theta_x-\frac{x}{c_1}(a+b+1-c_2)\theta_y \\
& +\frac{(2c_1+c_2-a-b-1)}{(1-x-y)(c_1+c_2-a-1)(c_1+c_2-b-1)}\Biggl([(1-x-y)^2-4xy]\theta_{xy} \\
& \qquad -y\left[(1-x-y)(a+b-c_2-2c_1+2)+2x(a+b-c_1+1)\right]\theta_x \\
& \qquad -\frac{x}{c_1}\left[2c_1y(a+b+1-c_2)+(1-x-y)(c_2-a-1)(c_2-b-1)\right]\theta_y \\
& \qquad -2abxy \qquad\qquad\qquad \Biggr)\Biggr\} F_4(a,b,c_1,c_2;x,y)\ .
\end{aligned}
\tag{75}
$$

The differential reduction for the parameters "b" and "c_2" follows from Eqs. (72), (74) and Eqs. (73), (75), respectively, and the symmetry property of the function F_4 :

(i) $F_4(a,b,c_1,c_2,c;x,y) = F_4(b,a,c_1,c_2,c;x,y)\ :\quad a \Leftrightarrow b\ ,$

(ii) $F_4(a,b,c_1,c_2,c;x,y) = F_4(a,b,c_2,c_1,c;y,x)\ :\quad c_1 \Leftrightarrow c_2\ ,\quad x \Leftrightarrow y\ .$

Table 2:

F_1	$\{a, b_1, b_2, c-a, c-b_1-b_2\} \in \mathbb{Z}$
F_2	$\{a, b_1, b_2, c_1-a, c_2-a, c_1+c_2-a, c_1-b_1, c_2-b_2\} \in \mathbb{Z}$
F_3	$\{a_1, a_2, b_1, b_2, c-a_1-a_2, c-b_1-b_2, c-a_2-b_1, c-a_1-b_2\} \in \mathbb{Z}$
F_4	$\{a, b, c_1-a, c_1-b, c_2-a, c_2-b, c_1+c_2-a, c_1+c_2-b\} \in \mathbb{Z}$

3.6 Appell hypergeometric functions: exceptional values of parameters

As was explained in Section 3.1, the differential reduction algorithm in application to Appell's functions could be written symbolically as follows:

$$R(x,y)F_1(\vec{A}+\vec{m};x,y) = [P_0(x,y) + P_1(x,y)\theta_x + P_2(x,y)\theta_y]\, F_1(\vec{A};x,y) \ , \tag{76}$$

$$S(x,y)F_j(\vec{A}+\vec{m};x,y) = [Q_0(x,y)+Q_1(x,y)\theta_x+Q_2(x,y)\theta_y+Q_3(x,y)\theta_x\theta_y]\, F_j(\vec{A};x,y) \ , \tag{77}$$

where $j = 2, 3, 4$, $\vec{m}$ is a set of integers, $\vec{A}$ is a set of parameters, R, S, P_i, Q_i are some polynomials and $\theta_x = x\partial_x(\theta_y = y\partial_y)$.

However, there is a special subset of values of parameters when the results of differential reduction, Eqs. (76) and (77), have a simpler form. This set of exceptional values of parameters can be defined from (i) condition that the hypergeometric function entering in the l.h.s. of Eqs. (41)-(43), (52)-(54), (63)-(65), (74), (75), is expressible in terms of simpler hypergeometric functions (e.g. Gauss hypergeometric functions); (ii), condition that some of the coefficients entering in the inverse differential relations are equal to zero(infinity).

For Appell's hypergeometric functions, F_1, F_2, F_3 and F_4, the exceptional set of parameters are enumerated in Table 2.

It is not surprising that the set of exceptional values of the parameters coincides with the set of parameters defining the condition of irreducibility of monodromy group of the corresponding hypergeometric functions (see [24] and references therein). We are expecting that the condition of irreducibility of the Mellin-Barnes integral [25] can be used as a criterion of irreducibility of Feynman Diagram.

4 pfq - differential reduction of hypergeometric function ${}_pF_{p-1}$.

4.1 Non-exceptional values of parameters

In this section, we will present the Mathematica[1] based package **pfq** for differential reduction of the hypergeometric function ${}_pF_{p-1}$. In contrast to the version presented in [22] the current version deals with non-exceptional and exceptional values of parameters. The Takayama algorithm is implemented in file "**pfq.m**", and an example of application is given in the file "example-pfq.m".

---

[1] It was tested for Mathematica 7.0.

The program could be loaded in a standard way:

<< "pfq.m"

and includes two routines: **ToGroebnerBasis**[...] and **explicitForm**[...].
The main routine

$$\textbf{ToGroebnerBasis}[\text{Argumentsvector}], \tag{78}$$

calculates explicitly the ratio of the functions $\{R_k\}$ and S of Eq. (19). The "Argumentsvector" in Eq. (78) is the set of parameters of the hypergeometric function in the l.h.s of Eq. (19) with an explicit set of integer numbers, by which each parameter should be shifted:

$$\text{Argumentsvector} = \{\{\vec{a} + \vec{M}\}, \{\vec{b} + \vec{K}\}, \{x\}\}\ , \tag{79}$$

where $\vec{M}, \vec{K}$ are integers, $\vec{a}, \vec{b}$ are any symbols, and x denotes the argument of the hypergeometric function. The output of **ToGroebnerBasis**[...] has the following structure:

$$\{\{Q_1, \cdots, Q_{p+1}\}, \{\{a_1, \cdots, a_{p+1}\}, \{1+b_1, \cdots, 1+b_p\}, x\}, \text{factor}\}, \tag{80}$$

where

- $\{\{a_1, \cdots, a_{p+1}\}, \{1+b_1, \ldots, 1+b_p\}, x\}$ are the set of parameters and the argument of the resulting hypergeometric function entering in r.h.s. of Eq. (19);
- $\{Q_1, \cdots, Q_{p+1}\}$ are the rational functions where the numbering corresponds to power of θ^i, $i = 0, \ldots, p$;
- factor is an overall factor.

The explicit form of Eqs. (78)-(80) is the following:

$$\begin{aligned}
&{}_{p+1}F_p\left(\begin{array}{c} a_1{+}M_1, \cdots, a_{p+1}{+}M_{p+1} \\ b_1{+}K_1, \cdots, b_p{+}K_p \end{array} \middle| x \right) \\
&= \text{factor} \times \left(Q_1{+}Q_2\theta{+}\cdots{+}Q_{p+1}\theta^p \right) {}_{p+1}F_p\left(\begin{array}{c} a_1, \cdots, a_{p+1} \\ 1{+}b_1, \cdots, 1{+}b_p \end{array} \middle| x \right) .
\end{aligned} \tag{81}$$

1. **Example 1** [2] : reduction of ${}_3F_2$
   ToGroebnerBasis[{{1+a$_1$,2+a$_2$, a$_3$},{1+b$_1$, b$_2$+2},x}]

   IntegerPart={1,2,0,1,2}   changeVector={-1,-2,0,0,-1}   ,

---

[2] All functions in the package HYPERDIRE generate output without additional simplification. This is done for the maximum efficiency of algorithm. To get a simpler form of output, we recommended to use additional **Simplify** command. In particular, all considered examples correspond to **FullSimplify**[**ToGroebnerBasis**[...]].

$$\left\{-\tfrac{(a_2-a_3+1)(b_2+1)}{(a_2+1)(a_3-b_2-1)}, -\tfrac{(b_2+1)\left(xa_2^2-(a_3x-x+b_1)a_2+xa_1(a_2-a_3+1)+b_1b_2\right)}{xa_1a_2(a_2+1)(a_3-b_2-1)}, -\tfrac{(b_2+1)(-a_3x+x+(x-1)a_2+b_2)}{xa_1a_2(a_2+1)(a_3-b_2-1)}\right\} ,$$

$$\left\{\left\{a_1, a_2, a_3\right\}, \left\{b_1+1, b_2+1\right\}, x\right\}, 1\}$$

corresponds to

$$\begin{aligned}
{}_3F_2\left(\left.\begin{array}{c} a_1+1, a_2+2, a_3 \\ b_1+1, b_2+2 \end{array}\right| z\right) &= \left[-\frac{(a_2-a_3+1)(b_2+1)}{(a_2+1)(a_3-b_2-1)}\right. \\
&-\frac{(b_2+1)\left(xa_2^2-(a_3x-x+b_1)a_2+xa_1(a_2-a_3+1)+b_1b_2\right)}{xa_1a_2(a_2+1)(a_3-b_2-1)}\theta \\
&\left.-\frac{(b_2+1)(-a_3x+x+(x-1)a_2+b_2)}{xa_1a_2(a_2+1)(a_3-b_2-1)}\theta^2\right] {}_3F_2\left(\left.\begin{array}{c} a_1, a_2, a_3 \\ b_1+1, b_2+1 \end{array}\right| z\right) . \qquad (82)
\end{aligned}$$

2. **Example 2**: reduction of ${}_4F_3$
   ToGroebnerBasis [{{1+a$_1$,1+a$_2$, a$_3$,a$_4$},{1+b$_1$, b$_2$+1,b$_3$},x}] ,

   IntegerPart={1,1,0,0,1,1,0} changeVector={-1,-1,0,0,0,0,1}

   $$\left\{1, \tfrac{1}{a_2}+\tfrac{1}{b_3}+\tfrac{1}{a_1}, \tfrac{a_1+a_2+b_3}{a_1a_2b_3}, \tfrac{1}{a_1a_2b_3}\right\}, \left\{\left\{a_1, a_2, a_3, a_4\right\}, \left\{b_1+1, b_2+1, b_3+1\right\}, x\right\}, 1\}$$

   corresponds to

$$\begin{aligned}
{}_4F_3\left(\left.\begin{array}{c} 1+a_1, 1+a_2, a_3, a_4 \\ 1+b_1, 1+b_2, b_3 \end{array}\right| z\right) &= \left[1+\left(\frac{1}{a_2}+\frac{1}{b_3}+\frac{1}{a_1}\right)\theta\right. \\
&\left.+\frac{a_1+a_2+b_3}{a_1a_2b_3}\theta^2+\frac{1}{a_1a_2b_3}\theta^3\right] {}_4F_3\left(\left.\begin{array}{c} a_1, a_2, a_3, a_4 \\ b_1+1, b_2+1, b_3+1 \end{array}\right| x\right) . \qquad (83)
\end{aligned}$$

In both these examples, IntegerPart $= \{\cdots\}$ means the set of integer values of parameters in the original hypergeometric function, e.g. IntegerPart $= \{1,2,0,1,2\}$ in `Example 1`, and changeVector $= \{\cdots\}$ corresponds to values of the parameters which should be changed, e.g. changeVector $= \{-1,-2,0,0,-1\}$ in `Example 2`.

Routine **explicitForm**[…] convert the results of reduction, Eq. (81), to Mathematica-standard expression for generalized hypergeometric function.

Example 3 : reduction of ${}_2F_1$.
answer=**ToGroebnerBasis** [{1+a_1,1+a_2},{1+b_1},x}];

IntegerPart={1,1,1} changeVector={-1,-1,0}

explicitForm[answer]

$$\frac{\text{HypergeometricPFQ}(\{a_1,a_2\},\{b_1+1\},x)}{1-x} + \frac{a_1 a_2 x (a_1+a_2-b_1)\,\text{HypergeometricPFQ}(\{a_1+1,a_2+1\},\{b_1+2\},x)}{(b_1+1)(a_1 a_2 - a_1 a_2 x)}$$

For non-exceptional values of the parameters the differential reduction is performed with the help of Eqs. (10),(11),(13),(14). The higher powers of the operators θ^k are expressed with the help of the differential equation for the hypergeometric function, Eq. (9). Also, the following relation is used in some cases:

$$ {}_pF_q\left(\begin{array}{c} \{a_i+m\}_p \\ \{b_k+m\}_q \end{array} \middle| z \right) = \frac{\prod_{k=1}^q \{(b_k)_m\}}{\prod_{j=1}^p \{(a_j)_m\}} \left(\frac{d}{dz} \right)^m {}_pF_q\left(\begin{array}{c} \{a_i\}_p \\ \{b_k\}_q \end{array} \middle| z \right) . \tag{84}$$

4.2 Exceptional values of parameters

When some of the upper parameters of the initial hypergeometric function are integer, then the higher power of the differential operator can be excluded with the help of Eq. (21). In this case, the input of **ToGroebnerBasis**[...] does not change, and the output of **ToGroebnerBasis**[...] has the following structure (case when only one upper parameter is integer):

$$\{\{\{Q_1,\cdots,Q_p\},\{\{1+a_1,\ldots,1+a_p\},\{2+b_1,\cdots,2+b_p\},x\},\text{factor1}\},\{Q_{p+1},\{\},\text{factor2}\}\}, \tag{85}$$

where

- $\{\{1+a_1,\cdots,1+a_{p+1}\},\{2+b_1,\ldots,2+b_p\},x\}$ are the set of parameters and argument of the resulting hypergeometric function;
- **if upper parameter is integer, the proper a_j is equal to zero**;
- $\{Q_1,\cdots,Q_p\}$ are the rational functions where the numbering corresponds to power of θ^i, $i=0,\ldots,p-1$;
- factor is overall factor;
- Q_{p+1} is the resulting polynomial entering in the r.h.s. of Eq. (21).

In the explicit form

$$\begin{aligned} & {}_{p+1}F_p\left(\begin{array}{c} M_1,\cdots,M_r,a_{r+1}+M_{r+1},\cdots,a_{p+1}+M_{p+1} \\ b_1+K_1,\cdots,b_p+K_p \end{array} \middle| x \right) \\ & = \text{factor1} \times \left(Q_1+Q_2\theta+\cdots+Q_p\theta^{p-1} \right) {}_{p+1}F_p\left(\begin{array}{c} 1,\cdots,1_r,1+a_{r+1},\cdots,1+a_{p+1} \\ 2+b_1,\cdots,2+b_p \end{array} \middle| x \right) \\ & +\text{factor2} \times Q_{p+1} \ , \end{aligned} \tag{86}$$

where $\vec{M}$ is a set of integers.

Example 4 : reduction of ${}_3F_2$ with an integer parameter.

ToGroebnerBasis [{{3,1+a_2,1+a_3},{2+b_1,2+b_2},x}];

IntegerPart={3,1,1,2,2}   changeVector={-2,0,0,0,0}

$$\left\{\left\{\left\{\frac{(b_1+1)(b_2+1)-x(a_2+1)(a_3+1)}{2(x-1)}+1,\frac{1}{2}\left(\frac{-x(a_2+a_3+2)+b_1+b_2+2}{x-1}+3\right)\right\},\right.\right.$$

$$\left.\left.\{\{1,a_2+1,a_3+1\},\{b_1+2,b_2+2\},x\},1\right\},\{-\frac{(b_1+1)(b_2+1)}{2(x-1)},\{\},1\}\right\},$$

In an explicit form it is

$$
\begin{aligned}
{}_3F_2\left(\left.\begin{array}{c}3,a_2{+}1,a_3{+}1\\ b_1{+}2,b_2{+}2\end{array}\right| x\right) &= \left[\frac{(b_1{+}1)\,(b_2{+}1){-}x\,(a_2{+}1)\,(a_3{+}1)}{2(x-1)}+1\right.\\
&+\left.\frac{1}{2}\left(\frac{-x\,(a_2+a_3+2)+b_1+b_2+2}{x-1}+3\right)\theta\right]{}_3F_2\left(\left.\begin{array}{c}1,a_2+1,a_3+1\\ b_1+2,b_2+2\end{array}\right| x\right)\\
&-\frac{(b_1+1)\,(b_2+1)}{2(x-1)}\,.
\end{aligned}
$$

Other useful relations which are implemented in the function **ToGroebnerBasis**[...] are the relations derived in [26]:

$$
\begin{aligned}
&{}_pF_q\left(\left.\begin{array}{c}b_1+m_1,\cdots,b_n+m_n,a_{n+1},\cdots,a_p\\ b_1,\cdots,b_n,b_{n+1},\cdots,b_q\end{array}\right| z\right)\\
&=\sum_{j_1=0}^{m_1}\cdots\sum_{j_n=0}^{m_n}A(j_1,\cdots j_n)z^{J_n}{}_{p-n}F_{q-n}\left(\left.\begin{array}{c}a_{n+1}+J_n,\cdots a_p+J_n\\ b_{n+1}+J_n,\cdots,b_q+J_n\end{array}\right| z\right)\;,
\end{aligned}
\tag{87}
$$

where m_j are positive integers, $J_n = j_1 + \cdots + j_n$, and

$$
A(j_1,\cdots j_n) \;=\; \binom{m_1}{j_1}\cdots\binom{m_n}{j_n}\frac{(b_2{+}m_2)_{J_1}(b_3{+}m_3)_{J_2}\cdots(b_n{+}m_n)_{J_{n-1}}(a_{n+1})_{J_n}\cdots(a_p)_{J_n}}{(b_1)_{J_1}(b_2)_{J_2}\cdots(b_n)_{J_n}(b_{n+1})_{J_n}\cdots(b_q)_{J_n}}\,.
\tag{88}
$$

Example 5 : reduction of ${}_3F_2$ with integer difference values of parameters.
ToGroebnerBasis [{{3+b_1,1+a_2,1+a_3},{2+b_1,2+b_2},x}];

IntegerPart={3,1,1,2,2}   changeVector={-1,-1,-1}

$$\{\left\{-\frac{b_2+1}{(x-1)(b_1+2)},-\frac{(a_2x+a_3x-b_1x-x+b_1-b_2+1)(b_2+1)}{(x-1)xa_2a_3(b_1+2)}\right\},\{\{a_2,a_3\},\{b_2+1\},x\},1\}$$

In an explicit form:

$$
\begin{aligned}
& {}_3F_2\left(\begin{array}{c} 3+b_1, 1+a_2, 1+a_3 \\ 2+b_1, 2+b_2 \end{array} \middle| \, x \right) \\
& = \left[-\frac{b_2+1}{(x-1)(b_1+2)} - \frac{(a_2x+a_3x-b_1x-x+b_1-b_2+1)(b_2+1)}{(x-1)xa_2a_3(b_1+2)}\theta \right] {}_2F_1\left(\begin{array}{c} a_2, a_3 \\ b_2+1 \end{array} \middle| \, x \right) .
\end{aligned}
\tag{89}
$$

The following relations:

$$
\theta^k {}_{p+1}F_p\left(\begin{array}{c} A, \vec{a} \\ 1+A, \vec{b} \end{array} \middle| \, z \right) = (-A)^k {}_{p+1}F_p\left(\begin{array}{c} A, \vec{a} \\ 1+A, \vec{b} \end{array} \middle| \, z \right) - \sum_{j=0}^{k-1} (-A)^{k-j} \theta^j {}_pF_{p-1}\left(\begin{array}{c} \vec{a} \\ \vec{b} \end{array} \middle| \, z \right) ,
$$

$$
\left(\frac{\theta}{A}\right)^q {}_{p+1}F_p\left(\begin{array}{c} \{A\}_r, \vec{a} \\ \{1+A\}_r, \vec{b} \end{array} \middle| \, z \right) = \sum_{j=0}^{q} (-1)^{(j+q)} \binom{q}{j} {}_{p+1-j}F_{p-j}\left(\begin{array}{c} \{A\}_{r-j}, \vec{a} \\ \{1+A\}_{r-j}, \vec{b} \end{array} \middle| \, z \right) ,
$$

$$
{}_pF_q\left(\begin{array}{c} 1, \{a_i\}_{p-1} \\ 2, \{b_k\}_{q-1} \end{array} \middle| \, z \right) = \frac{1}{z} \frac{\prod_{l=1}^{q-1}(b_l-1)}{\prod_{j=1}^{p-1}(a_j-1)} \left[{}_{p-1}F_{q-1}\left(\begin{array}{c} \{a_i-1\}_{p-1} \\ \{b_k-1\}_{q-1} \end{array} \middle| \, z \right) - 1 \right] ,
\tag{90}
$$

where $q \leq r$ and $a_j, b_k \neq 1$ are implemented in Mathematica 7.0.

5 AppellF1F4 - Mathematica based program for differential reduction of Appell's functions F_1, F_2, F_3, F_4

In this section, we will present the MATHEMATICA[3] based program **AppellF1F4** for differential reduction of the Appell hypergeometric functions F_1, F_2, F_3 and F_4. The program is available in [38]. The current version, 1.0, deals with non-exceptional values of a parameters only.

The program could be loaded in a standard way:

<< ”AppellF1F4.m”

The package includes the following basic routines:

$$
\begin{aligned}
& \textbf{F1IndexChange}[\text{changingVector}, \text{parameterVector}], && (91) \\
& \textbf{F2IndexChange}[\text{changingVector}, \text{parameterVector}], && (92) \\
& \textbf{F3IndexChange}[\text{changingVector}, \text{parameterVector}], && (93) \\
& \textbf{F4IndexChange}[\text{changingVector}, \text{parameterVector}]. && (94)
\end{aligned}
$$

and

$$
\textbf{explicitFormF1}[\ldots] . \tag{95}
$$

---

[3] It was tested for MATHEMATICA 7.0.

The "changingVector" in Eq. (91)-(94) is the set of integers at which we want to change the values of parameters of the Appell functions (vector $\vec{m}$ in Eqs. (76), (77)). The set of parameters of Appell's function are defined in the list "parameterVector" (corresponds to vector $\vec{A}+\vec{m}$ and arguments x, y in the l.h.s. of Eqs. (76), (77)). We want to point out that enumeration of parameters and arguments in the "parameterVector" correspond one-to-one to Eqs. (33), (44), (55), (67).

The output of **F1IndexChange**[] and **F{234}IndexChange**[], has a little different structure.

5.1 Appell function F_1

The structure of output of **F1IndexChange**[] is the following:

$$\{\{\mathrm{A,B,C}\}, \{\mathrm{parameterVectorNew}\}, \{\mathrm{AppellF1}\}\}, \tag{96}$$

where

1. parameterVectorNew is the set of new parameters of Appell function F_1 ;
2. A, B, C are the rational functions corresponding to the ratio of $P_0/R, P_1/R$ and P_2/R of functions entering in Eq. (76).

Example 6 [4]: reduction of F_1.

F1IndexChange[{1,–1,0,0}, {a,b_1,b_2,c,z_1,z_2}]

$$\left\{\left\{\frac{a(-\mathrm{z1})+a+\mathrm{b1}\mathrm{z}_1+\mathrm{b}_2\mathrm{z}_2-c-\mathrm{z}_1+1}{a-c+1}, -\frac{(\mathrm{z}_1-1)(a-\mathrm{b}_1+1)}{(\mathrm{b}_1-1)(a-c+1)}, \frac{\mathrm{z}_2-1}{a-c+1}\right\}, \{a+1, \mathrm{b}_1-1, \mathrm{b}_2, c, \mathrm{z}_1, \mathrm{z}_2\}, \mathrm{AppellF1}\right\}$$

In an explicit form:

$$\begin{aligned} &F_1(a, b_1, b_2, c; z_1, z_2) = \\ &\left[\frac{-az_1+a+b_1z_1+b_2z_2-c-z_1+1}{a-c+1} - \frac{(z_1-1)(a-b_1+1)}{(b_1-1)(a-c+1)}\theta_1 + \frac{z_2-1}{a-c+1}\theta_2\right] \\ &\times F_1(a+1, b_1-1, b_2, c; z_1, z_2). \end{aligned} \tag{97}$$

Routine **explicitFormF1**[$\cdots$] converts the result of reduction to Mathematica standard expression for the Appell function F_1.

Example 7 : reduction of F_1.
result = **F1IndexChange**[{1,–1,0,0}, {a,b_1,b_2,c,z$_1$,z$_2$}]

explicitFormF1[result]

---

[4] See footnote 2.

$$-\tfrac{(a+1)(z_1-1)z_1(a-b_1+1)}{c(a-c+1)}\text{AppellF1}\left(a+2;b_1,b_2;c+1;z_1,z_2\right)$$
$$+\tfrac{(a+1)b_2(z_2-1)z_2}{c(a-c+1)}\text{AppellF1}\left(a+2;b_1-1,b_2+1;c+1;z_1,z_2\right)$$
$$+\tfrac{(a(-z_1)+a+b_1z_1+b_2z_2-c-z_1+1)}{a-c+1}\text{AppellF1}\left(a+1;b_1-1,b_2;c;z_1,z_2\right)\ .$$

5.2 Appell functions F_2, F_3, F_4

The output of **F2IndexChange**[], **F3IndexChange**[], and **F4IndexChange**[], is similar and has the following structure:

$$\{\{\text{A,B,C,D}\},\{\text{parameterVectorNew}\},\{\text{NameOfFunction}\},\tag{98}$$

where

1. NameOfFunction is the name of Appell functions under reduction: AppellF2, AppellF3 or AppellF4 ;
2. parameterVectorNew is the set of new parameters of the Appell function;
3. A, B, C, D are the rational functions corresponding to the ratio of $Q_0/S, Q_1/S, Q_2/S$ and Q_3/S of functions entering in Eq. (77).

Example 8 : reduction of F_2.

F2IndexChange[{0,0,1,1,0}, {a,b$_1$,b$_2$,c$_1$,c$_2$,z$_1$,z$_2$}]

$$\left\{\left\{\tfrac{az_2(c_1(z_1-1)-b_1z_1)}{c_1(z_1-1)(b_2-c_2+1)}+1,\tfrac{1-\frac{z_2(a+z_1(b_1-c_1))}{(z_1-1)(b_2-c_2+1)}}{c_1},\tfrac{c_1(z_1-1)(z_2-1)-b_1z_1z_2}{c_1(z_1-1)(b_2-c_2+1)},-\tfrac{z_1+z_2-1}{c_1(z_1-1)(b_2-c_2+1)}\right\},\right.$$
$$\left.\{a,b_1,b_2+1,c_1+1,c_2,z_1,z_2\},\text{AppellF2}\right\}$$

In an explicit form:

$$\begin{aligned}&F_2(a,b_1,b_2,c_1,c_2,z_1,z_2)=\\&\left[\frac{az_2\left(c_1\left(z_1-1\right)-b_1z_1\right)}{c_1\left(z_1-1\right)\left(b_2-c_2+1\right)}+1+\frac{1-\frac{z_2(a+z_1(b_1-c_1))}{(z_1-1)(b_2-c_2+1)}}{c_1}\theta_1+\frac{c_1\left(z_1-1\right)\left(z_2-1\right)-b_1z_1z_2}{c_1\left(z_1-1\right)\left(b_2-c_2+1\right)}\theta_2\right.\\&\left.-\frac{z_1+z_2-1}{c_1\left(z_1-1\right)\left(b_2-c_2+1\right)}\theta_1\theta_2\right]F_2(a,b_1,b_2+1,c_1+1,c_2;z_1,z_2)\ .\end{aligned}\tag{99}$$

The Appell functions F_2, F_3 and F_4, are not built in the current version of Mathematica (version 7.0), so that the **AppellF1F4** package does not include a function similar to **explicitFormF1**[].

6 Application: reduction of Feynman Diagrams

6.1 Hypergeometric structure of Feynman Diagrams

The standard way of analytical evaluation of multiloop Feynman Diagrams is based on exploring of integration by parts relations [27] for the reduction of original diagram to set so-called master-integrals. This reduction can be performed in an arbitrary space-time dimension (before construction of ε-expansion). Now there are a few ways of computer realization of this algorithm [28]. In a series of papers [13, 14] it was pointed out that the hypergeometric representation of Feynman Diagram in conjunction with the differential reduction algorithm allow one to make reduction directly in terms of hypergeometric functions. In particular, it was formulated the conjecture about the structure of the hypergeometric representation of Feynman Diagram (see as review [29]).

Let us consider the hypergeometric representation of the Feynman diagram $\Phi(n,\vec{j};\vec{z})$,

$$\Phi(n,\vec{j};\vec{z}) = \sum_{a=1}^{k} S_a(n,\vec{j},\vec{z}) H_a(\vec{\beta}_a;\vec{\lambda}_a;\vec{\xi}) \ , \tag{100}$$

where $H_a(\vec{\beta}_a;\vec{\lambda}_a;\vec{\xi})$ are the Horn-type hypergeometric functions defined by Eq. (1), $\vec{j}$ is a list of the powers of the propagators in the Feynman diagram, n is the space-time dimension, $\vec{\xi}$ are the arguments of the hypergeometric functions which are related to the kinematic invariants of the Feynman diagram, $\{\beta_a, \lambda_a\}$ are the linear combinations of $\vec{j}$ and n with a polynomial coefficients, and S_a are the rational functions of the variables $\vec{z}$ with the coefficients depending on n and $\vec{j}$.

In general, the r.h.s. of Eq. (100) can include the different types of the hypergeometric functions (e.g. ${}_{p+1}F_p$ and ${}_pF_{p-1}$, where p is integer, are allowed). Our proposition is that: (i) regardless the type of functions in the r.h.s. of Eq. (100) the number of basis elements is the same (up to a module of rational functions) for each term of the r.h.s. of Eq. (100); (ii) the number h of nontrivial master integrals following from integration-by-parts relations which are not expressible in terms of gamma functions is then equal to the number of basis elements L for each term of the r.h.s. of Eq. (100).

As illustration of the differential reduction algorithm and our conjecture, let us consider the diagrams shown in Figs.1, 2 and 3.

6.2 V_{0002}

Let us start from consideration of the two-loop propagator diagram V_{0002} depicted in Fig. (1) with one massive line and three-massless lines:

$$V_{000M}(j_1,\sigma_2,\sigma_1,j_4;p^2) = \int \frac{d^n(k_1k_2)}{[(k_2-p)^2]^{j_1}[(k_1-k_2)^2]^{\sigma_1}[k_1^2]^{\sigma_2}[k_2^2-M^2]^{j_4}} \ . \tag{101}$$



Figure 1: Diagrams implemented in package **HYPERDIRE**. Bold and thin lines correspond to massive and massless propagators, respectively.

Such diagrams appear as one of the master-integrals in pole mass of heavy quarks. The Mellin-Barnes representation of Eq. (101) reads:

$$V_{000M}(j_1,\sigma_2,\sigma_1,j_4;p^2) = \frac{\left[i^{1-n}\pi^{\frac{n}{2}}\right]^2}{(p^2)^{j_1+\sigma_1+\sigma_2+j_4-n}} \frac{\Gamma(\frac{n}{2}-j_1)\Gamma(\frac{n}{2}-\sigma_1)\Gamma(\frac{n}{2}-\sigma_2)\Gamma(\sigma_1+\sigma_2-\frac{n}{2})}{\Gamma(j_1)\Gamma(\sigma_1)\Gamma(\sigma_2)\Gamma(j_4)\Gamma(n-\sigma_1-\sigma_2)}$$
$$\times \int dt \left(-\frac{M^2}{p^2}\right)^t \frac{\Gamma(-t)\Gamma(t+j_4)\Gamma(n-\sigma_1-\sigma_2-j_4-t)\Gamma(j_1+\sigma_1+\sigma_2+j_4-n+t)}{\Gamma\left(\sigma_1+\sigma_2+j_4-\frac{n}{2}+t\right)\Gamma\left(\frac{3n}{2}-j_1-\sigma_1-\sigma_2-j_4-t\right)} \,. \tag{102}$$

Closing the contour of integration in Eq. (102) on the right, we obtain

$$V_{000M}(j_1,\sigma_1,\sigma_2,j_4;p^2) = \frac{\left[i^{1-n}\pi^{\frac{n}{2}}\right]^2}{(p^2)^{j_1+\sigma_1+\sigma_2+j_4-n}} \frac{\Gamma(\frac{n}{2}-j_1)\Gamma(\frac{n}{2}-\sigma_1)\Gamma(\frac{n}{2}-\sigma_2)\Gamma(\sigma_1+\sigma_2-\frac{n}{2})}{\Gamma(j_1)\Gamma(\sigma_1)\Gamma(\sigma_2)\Gamma(j_4)\Gamma(n-\sigma_1-\sigma_2)}$$
$$\Biggl\{ \left(-\frac{p^2}{M^2}\right)^{j_4} \frac{\Gamma(j_4)\Gamma(n-\sigma_1-\sigma_2)\Gamma(j_1+\sigma_1+\sigma_2-n)}{\Gamma(\sigma_1+\sigma_2-\frac{n}{2})\Gamma(\frac{3n}{2}-j_1-\sigma_1-\sigma_2)}$$
$$\times\, {}_3F_2\left(\begin{array}{c} j_4,\; n-\sigma_1-\sigma_2,\; \frac{n}{2}+1-\sigma_1-\sigma_2 \\ n+1-j_1-\sigma_1-\sigma_2,\; \frac{3n}{2}-j_1-\sigma_1-\sigma_2 \end{array} \middle| \frac{p^2}{M^2} \right)$$
$$+ \left(-\frac{p^2}{M^2}\right)^{j_1+\sigma_1+\sigma_2+j_4-n} \frac{\Gamma(j_1)\Gamma(n-j_1-\sigma_1-\sigma_2)\Gamma(j_1+\sigma_1+\sigma_2+j_4-n)}{\Gamma(\frac{n}{2})\Gamma(\frac{n}{2}-j_1)}$$
$$\times\, {}_3F_2\left(\begin{array}{c} j_1,\; j_1+\sigma_1+\sigma_2+j_4-n,\; 1+j_1-\frac{n}{2} \\ 1+j_1+\sigma_1+\sigma_2-n,\; \frac{n}{2} \end{array} \middle| \frac{p^2}{M^2} \right) \Biggr\} \,. \tag{103}$$

Both hypergeometric functions in Eq. (103) are reducible to the ${}_2F_1$ functions with one unity upper parameter, namely,

$${}_2F_1\left(\begin{array}{c} 1, \frac{n}{2}+I_1 \\ \frac{3n}{2}+I_2 \end{array} \middle| z \right) \,, \quad {}_2F_1\left(\begin{array}{c} 1, I_1-\frac{n}{2} \\ I_2+\frac{n}{2} \end{array} \middle| z \right) \,, \tag{104}$$

respectively, plus rational functions of z. In accordance with Section 6.1, there is only one nontrivial master-integral in the framework of integration-by-parts relations and a few integrals expressible in terms of the product of Γ-functions.



Figure 2: Diagrams implemented in the package **HYPERDIRE**. Bold and thin lines correspond to massive and massless propagators, respectively.

In accordance with Tarasov's algorithm [30] there is one master integral of the same topology and two integrals expressible in terms of the product of Γ-functions [31].

For completeness, we will present a few first terms of ε-expansion of the master-integral. For $n = 4 - 2\varepsilon$, where ε is the parameter of dimensional regularization [32] and $j_1 = \sigma_1 = \sigma_1 = \sigma_2 = 1$, we have

$$\frac{V_{000M}(1,1,1,1;m^2)}{(i\pi^{2-\varepsilon})^2} = \frac{1}{2\varepsilon}(-p^2)^{-2\varepsilon}\frac{\Gamma^3(1-\varepsilon)\Gamma(1+2\varepsilon)}{(1-2\varepsilon)\Gamma(3-3\varepsilon)}\frac{p^2}{M^2}$$
$$\left\{ {}_2F_1\left(\begin{array}{c} 1,\ 1-\varepsilon \\ 3-3\varepsilon \end{array} \middle| \frac{p^2}{M^2} \right) - \frac{1}{\varepsilon}\left(\frac{-p^2}{M^2}\right)^{-1+2\varepsilon} \frac{\Gamma(1+\varepsilon)\Gamma(3-3\varepsilon)}{\Gamma(1-\varepsilon)\Gamma(2-\varepsilon)}\, {}_2F_1\left(\begin{array}{c} 1,\ \varepsilon \\ 2-\varepsilon \end{array} \middle| \frac{p^2}{M^2} \right) \right\}, \tag{105}$$

where we use $i^{-n}(p^2)^{n/2} = (-p^2)^{n/2}$. For the second ${}_2F_1$ hypergeometric function the all-order



Figure 3: Diagrams implemented in the the package **HYPERDIRE**. Bold and thin lines correspond to massive and massless propagators, respectively.

ε-expansion are available:

$$ {}_2F_1\left(\begin{array}{c} 1,\ \varepsilon \\ 2-\varepsilon \end{array} \middle| z \right) = \frac{1-\varepsilon}{2(1-2\varepsilon)z} \left\{ 1+z-(1-z)^{1-2\varepsilon} - 2(1-z)^{1-2\varepsilon} \varepsilon \sum_{j=1}^{\infty} \varepsilon^j \sum_{k=1}^{j} (-2)^{j-k} S_{k,j-k+1}(z) \right\}, \tag{106} $$

where $\mathrm{Li}_n(z)$ and $\mathrm{S}_{a,b}(z)$ are classical and Nielsen polylogartihms, respectively. For the first hypergeometric function we apply the following relation:

$$ \frac{(1-2\varepsilon)}{2-3\varepsilon} z\, {}_2F_1\left(\begin{array}{c} 1,\ 1-\varepsilon \\ 3-3\varepsilon \end{array} \middle| z \right) = 1-(1-z)\, {}_2F_1\left(\begin{array}{c} 1,\ 1-\varepsilon \\ 2-3\varepsilon \end{array} \middle| z \right) \ . \tag{107} $$

The first coefficients of the ε expansion read:

$$
\begin{aligned}
&(-p^2)^{2\varepsilon}(1-2\varepsilon)^2\frac{V_{000M}(1,1,1,1;m^2)}{(i\pi^{2-\varepsilon})^2\Gamma^2(1+\varepsilon)} = \frac{1}{2\varepsilon^2}\frac{\Gamma^3(1-\varepsilon)\Gamma(1+2\varepsilon)}{\Gamma^2(1+\varepsilon)\Gamma(1-3\varepsilon)}\Bigg[1 \\
&+\varepsilon\Bigg\{1+2\frac{1-z}{z}\ln(1-z)+2\ln(-z)\Bigg\} \\
&+\varepsilon^2\Bigg\{3+2\ln^2(-z)+4\zeta_2-2\frac{1-z}{z}\left[\ln^2(1-z)-\ln(-z)\ln(1-z)+2\mathrm{Li}_2\left(z\right)\right]\Bigg\} \\
&+\varepsilon^3\Bigg\{9+\frac{4}{3}\ln^3(-z)+8\zeta_3+8\zeta_2\ln(-z) \\
&\qquad +2\frac{1-z}{z}\Bigg[4\mathrm{S}_{1,2}(z)+\frac{2}{3}\ln^3(1-z)+4\ln(1-z)\mathrm{Li}_2\left(z\right)-5\mathrm{Li}_3\left(z\right) \\
&\qquad\quad +2\zeta_2\ln(1-z)-\ln(-z)\mathrm{Li}_2\left(z\right)-\ln(-z)\ln^2(1-z)+\ln^2(-z)\ln(1-z)\Bigg]\Bigg]\Bigg\} \\
&+\varepsilon^4\Bigg\{27+\frac{2}{3}\ln^4(-z)+40\zeta_4+8\zeta_2\ln^2(-z)+16\zeta_3\ln(-z) \\
&\qquad -2\frac{1-z}{z}\Bigg[8\mathrm{S}_{1,3}(z)-10\mathrm{S}_{2,2}(z)+14\mathrm{Li}_4\left(z\right)+\frac{1}{3}\ln^4(1-z)+2\zeta_2\mathrm{Li}_2\left(z\right) \\
&\qquad\quad +8\ln(1-z)\mathrm{S}_{1,2}(z)-10\ln(1-z)\mathrm{Li}_3\left(z\right)-4\zeta_3\ln(1-z) \\
&\qquad\quad +4\ln^2(1-z)\mathrm{Li}_2\left(z\right)+2\zeta_2\ln^2(1-z)+\ln^2(-z)\ln^2(1-z) \\
&\qquad\quad -2\ln(-z)\mathrm{S}_{1,2}(z)+\ln(-z)\mathrm{Li}_3\left(z\right)-\frac{2}{3}\ln(-z)\ln^3(1-z) \\
&\qquad\quad -4\zeta_2\ln(-z)\ln(1-z)-2\ln(-z)\ln(1-z)\mathrm{Li}_2\left(z\right) \\
&\qquad\quad +\ln^2(-z)\mathrm{Li}_2\left(z\right)-\frac{2}{3}\ln^3(-z)\ln(1-z)\Bigg]\Bigg]\Bigg\}+\mathcal{O}(\varepsilon^5)\Bigg]\ , \qquad (108)
\end{aligned}
$$

where $z = p^2/M^2$.

To check Eq. (108), we evaluate the first few coefficients of the ε expansion of the original diagram in the large-mass limit [33] using the program packages developed in Refs. [34] to find agreement.

6.3 E_{10002}

Let us consider the 3-loop bubble diagram E_{10002} depicted in Fig. (1) with two massive lines and three massless ones:

$$
E_{10002}(\lambda,\alpha_1,\alpha_2,\beta,\sigma,m^2,M^2) = \int \frac{d^n(k_1k_2k_3)}{[(k_1-k_3)^2]^{\alpha_1}[k_1^2]^{\alpha_2}[(k_2-k_3)^2]^{\beta}[k_3^2-m^2]^{\lambda}[k_2^2-M^2]^{\sigma}}\ . \qquad (109)
$$

Its Mellin-Barnes representation reads:

$$E_{10002}(\lambda,\alpha_1,\alpha_2,\beta,\sigma,m^2,M^2) = \left[i^{1-n}\pi^{\frac{n}{2}}\right]^3 (-m^2)^{n-a_{1,2,\lambda}}(-M^2)^{n/2-a_{\sigma,\beta}}$$
$$\times\frac{\Gamma\left(\frac{n}{2}-\alpha_1\right)\Gamma\left(\frac{n}{2}-\alpha_2\right)\Gamma\left(\frac{n}{2}-\beta\right)\Gamma\left(a_{1,2}-\frac{n}{2}\right)}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma(\sigma)\Gamma(\lambda)\Gamma(\beta)\Gamma\left(\frac{n}{2}\right)\Gamma(n-a_{1,2})}$$
$$\times\int dt \left(\frac{m^2}{M^2}\right)^t \frac{\Gamma(-t)\Gamma(\beta+t)\Gamma\left(a_{\beta,\sigma}-\frac{n}{2}+t\right)\Gamma\left(n-a_{1,2}+t\right)\Gamma\left(a_{1,2,\lambda}-n-t\right)}{\Gamma\left(t+\frac{n}{2}\right)}\ , \tag{110}$$

where

$$a_{\sigma,\beta} = \sigma+\beta\ , \quad a_{1,2,\lambda} = \alpha_1+\alpha_2+\lambda\ ,$$
$$a_{1,2,\beta,\lambda,\sigma} = \alpha_1+\alpha_2+\lambda+\beta+\sigma\ , \quad a_{1,2,\lambda,\beta} = \alpha_1+\alpha_2+\lambda+\beta\ . \tag{111}$$

Closing the contour of integration in Eq. (110) on the right, we obtain

$$E_{10002}(\lambda,\alpha_1,\alpha_2,\beta,\sigma,m^2,M^2) = \left[i^{1-n}\pi^{\frac{n}{2}}\right]^3 (-m^2)^{n-a_{1,2,\lambda}}(-M^2)^{n/2-a_{\sigma,\beta}}$$
$$\times\frac{\Gamma\left(\frac{n}{2}-\alpha_1\right)\Gamma\left(\frac{n}{2}-\alpha_2\right)\Gamma\left(\frac{n}{2}-\beta\right)\Gamma\left(a_{1,2}-\frac{n}{2}\right)}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma(\sigma)\Gamma(\lambda)\Gamma(\beta)\Gamma\left(\frac{n}{2}\right)\Gamma(n-a_{1,2})}$$
$$\Bigg[\frac{\Gamma(\beta)\Gamma\left(a_{\sigma,\beta}-\frac{n}{2}\right)\Gamma(n-a_{1,2})\Gamma(a_{1,2,\lambda}-n)}{\Gamma\left(\frac{n}{2}\right)}{}_3F_2\left(\left.\begin{array}{c}\beta, a_{\beta,\sigma}-\frac{n}{2}, n-a_{1,2}\\ \frac{n}{2}, 1+n-a_{1,2,\lambda}\end{array}\right| \frac{m^2}{M^2}\right)$$
$$+\left(\frac{m^2}{M^2}\right)^{a_{1,2,\lambda}-n}\frac{\Gamma(\lambda)\Gamma\left(a_{1,2,\lambda,\beta}-n\right)\Gamma\left(a_{1,2,\beta,\lambda,\sigma}-\frac{3n}{2}\right)\Gamma(n-a_{1,2,\lambda})}{\Gamma\left(a_{1,2,\lambda}-\frac{n}{2}\right)}$$
$$\times{}_3F_2\left(\left.\begin{array}{c}\lambda, a_{1,2,\lambda,\beta}-n, a_{1,2,\lambda,\beta,\sigma}-\frac{3n}{2}\\ a_{1,2,\lambda}-\frac{n}{2}, 1+a_{1,2,\lambda}-n\end{array}\right| \frac{m^2}{M^2}\right)\Bigg]\ . \tag{112}$$

Both hypergeometric functions in Eq. (112) are reducible to the ${}_2F_1$ functions with one unity upper parameter, namely,

$${}_2F_1\left(\left.\begin{array}{c}1, I_1-\frac{n}{2}\\ \frac{n}{2}+I_2\end{array}\right| z\right)\ , \quad {}_2F_1\left(\left.\begin{array}{c}1, I_1-\frac{3n}{2}\\ I_2-1\frac{n}{2}\end{array}\right| z\right)\ , \tag{113}$$

respectively, plus rational functions of z. For this diagram there is only one nontrivial master-integral in the framework of integration-by-parts relations and an integral expressible in terms of the product of Γ-functions.

For completeness, we will present a few first terms of the ε-expansion of the master-integral. For $n = 4 - 2\varepsilon$, where ε is the parameter of dimensional regularization [32] and $\alpha_1 = \alpha_2 = \sigma = \lambda = \beta = 1$, we have

$$E_{10002}(1,1,1,1,1,m^2,M^2) = \left[i\pi^{2-\varepsilon}\right]^3 (m^2)^{1-2\varepsilon}(M^2)^{-\varepsilon}$$
$$\times\frac{\Gamma(1-\varepsilon)\Gamma^2(1+\varepsilon)\Gamma(1+2\varepsilon)}{2\varepsilon^3(1-\varepsilon)^2(1-2\varepsilon)}\Bigg[{}_2F_1\left(\left.\begin{array}{c}1,\varepsilon\\ 2-\varepsilon\end{array}\right| \frac{m^2}{M^2}\right)$$
$$+\left(\frac{m^2}{M^2}\right)^{-1+2\varepsilon}\frac{\Gamma(2-\varepsilon)\Gamma(1+3\varepsilon)}{3(1-3\varepsilon)\Gamma^2(1+\varepsilon)}{}_2F_1\left(\left.\begin{array}{c}1,-1+3\varepsilon\\ 1+\varepsilon\end{array}\right| \frac{m^2}{M^2}\right)\Bigg]\ . \tag{114}$$

The all-order ε-expansion for the first hypergeometric function is given by Eq. (106) and for the second hypergeometric function we apply the following relation:

$$
{}_2F_1\left(\left. \begin{array}{c} 1, -1+3\varepsilon \\ 1+\varepsilon \end{array} \right| z \right) = 1 - \frac{1-3\varepsilon}{1-2\varepsilon} z + \frac{3\varepsilon(1-3\varepsilon)z(1-z)}{(1+\varepsilon)(1-2\varepsilon)} {}_2F_1\left(\left. \begin{array}{c} 1, 1+3\varepsilon \\ 2+\varepsilon \end{array} \right| z \right) . \qquad (115)
$$

The first coefficients of the ε expansion read:

$$
\begin{aligned}
& \frac{E_{10002}(1,1,1,1,1,m^2,M^2)}{\left[i\pi^{2-\varepsilon}\right]^3 \Gamma^3(1+\varepsilon)} (m^2)^{2\varepsilon} (M^2)^{-1+\varepsilon} = \frac{\Gamma(1-\varepsilon)\Gamma(1+2\varepsilon) z}{(1-\varepsilon)(1-2\varepsilon)^2 \Gamma(1+\varepsilon)} \\
& \times \Bigg[\frac{1}{6\varepsilon^3}(1+2z) + \frac{1}{6\varepsilon^2}\left[1 + 2(1-z)\ln z\right] \\
& + \frac{1}{3\varepsilon}\left[\frac{3}{2} + \ln z + (1-4z)\zeta_2 + (1-z)\left(\ln^2 z - 3\ln z \ln(1-z) - 3\mathrm{Li}_2(z)\right)\right] \\
& + \frac{3}{2} + \frac{1}{3}\zeta_2 + 3\zeta_3 + \ln z + \frac{1}{3}\ln^2 z - (1-z)\Bigg\{ \frac{10}{3}\zeta_3 - \frac{2}{9}\ln^3 z - 2\mathrm{S}_{1,2}(z) - \frac{2}{3}\zeta_2 \ln z \\
& \quad + \ln(1-z)\left[2\zeta_2 - 2\mathrm{Li}_2(z) + \ln^2 z\right] - \ln z \ln^2(1-z) + \ln z \mathrm{Li}_2(z) \Bigg\} \\
& + \varepsilon \Bigg\{ \frac{9}{2} + \zeta_2 - \frac{1}{3}\zeta_3 - \frac{11}{2}\zeta_4 + \frac{2}{9}\ln^3 z + \ln^2 z + \frac{1}{3}(9 + 2\zeta_2)\ln z \\
& \quad + (1-z)\Bigg\{ 7\zeta_4 - \mathrm{Li}_4(z) + \ln z \left[2\mathrm{S}_{1,2}(z) + \mathrm{Li}_3(z) - \frac{2}{3}\zeta_3 \right] \\
& \qquad + \frac{1}{9}\ln^4 z - 4\mathrm{S}_{1,3}(z) + \ln(1-z)\left[4\zeta_3 - 4\mathrm{S}_{1,2}(z) - \frac{2}{3}\ln^3 z \right] \\
& \qquad + 2\ln(1-z)\Big[\ln(1-z) - \ln z \Big]\Big[\zeta_2 - \mathrm{Li}_2(z) \Big] + \ln^2 z \ln^2(1-z) \\
& \qquad - \frac{2}{3}\ln z \ln^3(1-z) - \ln^2 z \left[\mathrm{Li}_2(z) - \frac{2}{3}\zeta_2 \right] \Bigg\} \Bigg\} \Bigg] , \qquad (116)
\end{aligned}
$$

where $z = m^2/M^2$. For $z = 1$, Eq.(116) coincides with the results of [35].

6.4 $B^q_{1122000}$ & B^q_{112200} & B^q_{11220} & B_{1122}

Let us consider the q-loop bubble type diagram $B^q_{1122000}$ [5] (see Fig. 2) defined as

$$\begin{aligned}
&B^q_{1122000}(m^2, M^2, \alpha_1, \alpha_2, \beta_1, \beta_2, \sigma_1, \cdots, \sigma_x, \rho_1, \cdots, \rho_r, \nu_1, \cdots, \nu_z) \\
&= \int \frac{d^n(k_1 \cdots k_q)}{[k_1^2]^{\rho_1} \cdots [k_{r-1}^2]^{\rho_{r-1}} [(k_1 + \cdots + k_{r-1} + P)^2]^{\rho_r}} \\
&\times \frac{1}{[k_r^2]^{\sigma_1} \cdots [k_{r+x-1}^2]^{\sigma_x} [k_{r+x}^2 - m^2]^{\alpha_1} [(k_r + \cdots + k_{r+x} + P)^2 - m^2]^{\alpha_2}} \\
&\times \frac{1}{[k_{r+x+1}^2]^{\nu_1} \cdots [k_{r+x+s}^2]^{\nu_s} [k_{q-1}^2 - M^2]^{\beta_1} [k_q^2 - M^2]^{\beta_2}} \ ,
\end{aligned} \tag{117}$$

where $P = k_{r+x+1} + \cdots + k_q$. and r, s and x are the number of massless lines. The Mellin-Barnes representation for diagram $B^q_{1122000}$ is:

$$\begin{aligned}
&B^q_{1122000}(m^2, M^2, \alpha_1, \alpha_2, \beta_1, \beta_2, \sigma_1, \cdots, \sigma_x, \rho_1, \cdots, \rho_r, \nu_1, \cdots, \nu_s) \\
&= \frac{\left[i^{1-n} \pi^{\frac{n}{2}} \right]^q (-m^2)^{\frac{n}{2} - \alpha_{1,2}} (-M^2)^{\frac{n}{2}(q-1) - a_{\rho,\sigma,\nu,\beta_{1,2}}}}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma(\beta_1)\Gamma(\beta_2)\Gamma\left(\frac{n}{2}\right)} \\
&\times \left\{ \Pi_{i=1}^x \frac{\Gamma(\frac{n}{2} - \sigma_i)}{\Gamma(\sigma_i)} \right\} \left\{ \Pi_{j=1}^r \frac{\Gamma(\frac{n}{2} - \rho_j)}{\Gamma(\rho_j)} \right\} \left\{ \Pi_{k=1}^s \frac{\Gamma(\frac{n}{2} - \nu_k)}{\Gamma(\nu_k)} \right\} \frac{\Gamma\left(\rho - \frac{n}{2}(r-1)\right)}{\Gamma\left(\frac{n}{2}r - \rho\right)} \\
&\times \int dt \left(\frac{M^2}{m^2} \right)^t \Gamma\left(\sigma - \tfrac{n}{2}x - t\right) \Gamma\left(a_{\beta_1,\beta_2,\nu,\rho,\sigma} - \tfrac{n}{2}(q-1) - t\right) \\
&\times \frac{\Gamma(\alpha_1 + t)\Gamma(\alpha_2 + t)\Gamma(\alpha_{1,2} - \frac{n}{2} + t)\Gamma(\frac{n}{2} + t)\Gamma(\frac{n}{2}(r+x) - a_{\rho,\sigma} + t)}{\Gamma(\alpha_{1,2} + 2t)\Gamma(\frac{n}{2}(x+1) - \sigma + t)} \\
&\times \frac{\Gamma(a_{\nu,\rho,\sigma} - \frac{n}{2}(q-3) - t)\Gamma(a_{\beta_1,\nu,\rho,\sigma} - \frac{n}{2}(q-2) - t)\Gamma(a_{\beta_2,\nu,\rho,\sigma} - \frac{n}{2}(q-2) - t)}{\Gamma(a_{\rho,\sigma} - \frac{n}{2}(r+x-1) - t)\Gamma(\beta_{1,2} + 2a_{\sigma,\rho,\nu} - n(q-2) - 2t)} \ ,
\end{aligned} \tag{118}$$

where $x + r + s = q - 2$, $q \geq 3$, and we have introduced the short notation,

$$\begin{aligned}
&\rho = \sum_{a=1}^r \rho_a \ , \quad \sigma = \sum_{a=1}^x \sigma_a \ , \quad \nu = \sum_{a=1}^s \nu_a \ , \quad a_{\beta_1,\beta_2,\nu,\rho,\sigma} = \beta_1 + \beta_2 + \sigma + \nu + \rho \ , \\
&\alpha_{1,2} = \alpha_1 + \alpha_2 \ , \quad \beta_{1,2} = \beta_1 + \beta_2 \ , \quad a_{\nu,\rho,\sigma} = \sigma + \nu + \rho \ , \quad a_{\rho,\sigma} = \sigma + \rho \ , \\
&a_{\beta_1,\nu,\rho,\sigma} = \beta_1 + \sigma + \nu + \rho \ , \quad a_{\beta_2,\nu,\rho,\sigma} = \beta_2 + \sigma + \nu + \rho \ .
\end{aligned} \tag{119}$$

The result could be written as a sum of 5 hypergeometric functions of type ${}_8F_7$ of argument m^2/M^2 or M^2/m^2. The expression is really long to be considered here.

More interesting particular cases could be derived when q, r, s, x take special values.

Diagram B^q_{112200} in Fig. 2 emerges from $B^q_{1122000}$ with $s = 0$, $\left\{ \Pi_{k=1}^s \frac{\Gamma(\frac{n}{2} - \nu_k)}{\Gamma(\nu_k)} \right\} = 1$ and $x + r = q - 2$. In this case, $\nu = 0$ and Eq. (118) is simplified due to $\lim_{\nu \to 0} \frac{\Gamma(a_{\nu,\rho,\sigma} - \frac{n}{2}(q-3) - s)}{\Gamma(a_{\rho,\sigma} - \frac{n}{2}(q-3) - s)} = 1$.

---

[5] We are indebted to A. Grozin who suggest us apply our technology to analyze diagram of this type.

The detailed analysis of this diagram was presented in [13]. The result for diagram B^q_{112200} can be expressed as a sum of four hypergeometric functions of type ${}_7F_6$ or of lower order function. The general result again is to be published. We present only the results of differential reduction. In the case when the powers of the propagators are all integers, the appearing hypergeometric functions are reducible to the following hypergeometric functions and their derivatives:[6]

$$\begin{array}{l}
\{1,\theta,\theta^2,\theta^3\} \times \; {}_4F_3\left(\left. \begin{array}{c} I_1-\frac{n}{2}(x-1), I_2-\frac{n}{2}x, I_3-\frac{n}{2}(x+1), \frac{1}{2}+I_4+\frac{n}{2}(q-x-2) \\ I_5+\frac{n}{2}, I_6+\frac{n}{2}(q-x-1), \frac{1}{2}+I_7-\frac{n}{2}x \end{array} \right| z \right) \\
\{1,\theta,\theta^2,\theta^3\} \times \; {}_4F_3\left(\left. \begin{array}{c} \frac{1}{2}+I_1-\frac{n}{2}, I_2-\frac{n}{2}(q-2), I_3-\frac{n}{2}(q-1), I_4-\frac{n}{2}q \\ I_5-\frac{n}{2}(q-x-1), I_6-\frac{n}{2}(q-x-2), \frac{1}{2}+I_7-\frac{n}{2}(q-1) \end{array} \right| z \right) \\
\{1,\theta,\theta^2,\theta^3\} \times \; {}_5F_4\left(\left. \begin{array}{c} 1, \frac{1}{2}+I_1, I_2-\frac{n}{2}(q-1), I_3-\frac{n}{2}(q-2), I_4-\frac{n}{2}(q-3) \\ I_5+\frac{n}{2}, \frac{1}{2}+I_6-\frac{n}{2}(q-2), I_7-\frac{n}{2}(q-x-2), I_8-\frac{n}{2}(q-x-3) \end{array} \right| z \right) \; ,
\end{array} \tag{120}$$

where I_k are arbitrary integers. In the last expression, some rational functions are also generated. In accordance with Section 6.1, this integral has four master-integrals for arbitrary values of x and q. Further simplification is possible for (i) $x = 0$ and (ii) $x = 1$. In these cases we have:

- for $x = 0$,

$$\begin{array}{l}
\{1,\theta\} \times \; {}_3F_2\left(\left. \begin{array}{c} 1, I_2-\frac{n}{2}, \frac{1}{2}+I_3+\frac{n}{2}(q-2) \\ I_4+\frac{n}{2}(q-1), \frac{1}{2}+I_5 \end{array} \right| z \right) \; , \\
\{1,\theta\} \times \; {}_2F_1\left(\left. \begin{array}{c} I_1+\frac{1}{2}-\frac{n}{2}, I_2-\frac{n}{2}q \\ I_3+\frac{1}{2}-\frac{n}{2}(q-1) \end{array} \right| z \right) \; , \\
\{1,\theta\} \times \; {}_3F_2\left(\left. \begin{array}{c} 1, \frac{1}{2}+I_1, I_2-\frac{n}{2}(q-1) \\ I_3+\frac{n}{2}.I_4+\frac{1}{2}-\frac{n}{2}(q-2) \end{array} \right| z \right) \; ,
\end{array} \tag{121}$$

  and in accordance with Section 6.1, this integral has two master-integrals only.

- for $x = 1$

$$\begin{array}{l}
\{1,\theta,\theta^2\} \times \; {}_4F_3\left(\left. \begin{array}{c} 1, I_1-\frac{n}{2}, I_2-n, \frac{1}{2}+I_3+\frac{n}{2}(q-3) \\ I_4+\frac{n}{2}, I_5+\frac{n}{2}(q-2), \frac{1}{2}+I_6-\frac{n}{2} \end{array} \right| z \right) \; , \\
\{1,\theta,\theta^2\} \times \; {}_3F_2\left(\left. \begin{array}{c} I_1+\frac{1}{2}-\frac{n}{2}, I_2-\frac{n}{2}(q-1), I_3-\frac{n}{2}q \\ I_4-\frac{n}{2}(q-3), I_5+\frac{1}{2}+\frac{n}{2}(q-1) \end{array} \right| z \right) \; , \\
\{1,\theta,\theta^2\} \times \; {}_4F_3\left(\left. \begin{array}{c} 1, \frac{1}{2}+I_1, I_2-\frac{n}{2}(q-1), I_3-\frac{n}{2}(q-2) \\ I_4+\frac{n}{2}, I_5+\frac{1}{2}-\frac{n}{2}(q-2), I_6+\frac{n}{2}(q-4) \end{array} \right| z \right) \; ,
\end{array} \tag{122}$$

and in accordance with Section 6.1, this integral has three nontrivial master-integrals only.

For a next diagram, B^q_{11220} (see Fig. 1), the Mellin-Barnes representation follows from Eq. (118) putting $r = 1, \rho = \rho_1 \to 0, z = 0$ and $x = q - 3$. For simplicity, we will present it an explicit form:

---

[6] Two of the original hypergeometric functions have a similar parameter structure and the same basis functions.

$$
\begin{aligned}
& B^{q}_{11220}(m^2, M^2, \alpha_1, \alpha_2, \beta_1, \beta_2, \sigma_1, \cdots, \sigma_{q-3}) \\
& = \frac{\left[i^{1-n}\pi^{\frac{n}{2}}\right]^q (-m^2)^{\frac{n}{2}-\alpha_{1,2}} (-M^2)^{\frac{n}{2}(q-1)-a_{\sigma,\beta_{1,2}}}}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma(\beta_1)\Gamma(\beta_2)\Gamma\left(\frac{n}{2}\right)} \left\{ \Pi_{i=1}^{q-3} \frac{\Gamma(\frac{n}{2}-\sigma_i)}{\Gamma(\sigma_i)} \right\} \\
& \times \int dt \left(\frac{M^2}{m^2}\right)^t \Gamma\left(\sigma - \tfrac{n}{2}(q-3) - t\right) \Gamma\left(a_{\beta_1,\beta_2,\sigma} - \tfrac{n}{2}(q-1) - t\right) \Gamma\left(\tfrac{n}{2} + t\right) \\
& \times \frac{\Gamma(\alpha_1+t)\Gamma(\alpha_2+t)\Gamma\left(\alpha_{1,2} - \frac{n}{2} + t\right) \Gamma\left(a_{\beta_1,\sigma} - \frac{n}{2}(q-2) - t\right) \Gamma\left(a_{\beta_2,\sigma} - \frac{n}{2}(q-2) - t\right)}{\Gamma(\alpha_{1,2}+2t)\Gamma\left(\beta_{1,2} + 2\sigma - n(q-2) - 2t\right)} \ . \qquad (123)
\end{aligned}
$$

and in agreement with Eq. (119) we have introduced the short notation,

$$
a_{\beta_1,\beta_2,\sigma} = \beta_1 + \beta_2 + \sigma \ , \quad a_{\beta_1,\sigma} = \beta_1 + \sigma \ , \quad a_{\beta_2,\sigma} = \beta_2 + \sigma \ . \qquad (124)
$$

The result could be written as a sum of four hypergeometric functions of type ${}_6F_5$ of the argument m^2/M^2 or M^2/m^2. Again result is to long to be published. The results of differential reduction follows from Eq. (120) by substitution $x = q - 3$:

$$
\begin{aligned}
& \{1, \theta, \theta^2, \theta^3\} \times {}_4F_3 \left(\begin{array}{c} I_1 - \frac{n}{2}(q-4), I_2 - \frac{n}{2}(q-3), I_3 - \frac{n}{2}(q-2), \frac{1}{2} + I_4 + \frac{n}{2} \\ I_5 + \frac{n}{2}, I_6 + n, \frac{1}{2} + I_7 - \frac{n}{2}(q-3) \end{array} \middle| z \right) \\
& \{1, \theta, \theta^2, \theta^3\} \times {}_4F_3 \left(\begin{array}{c} \frac{1}{2} + I_1 - \frac{n}{2}, I_2 - \frac{n}{2}(q-2), I_3 - \frac{n}{2}(q-1), I_4 - \frac{n}{2}q \\ I_5 - n, I_6 - \frac{n}{2}, \frac{1}{2} + I_7 - \frac{n}{2}(q-1) \end{array} \middle| z \right) \\
& \{1, \theta, \theta^2, \theta^3\} \times {}_4F_3 \left(\begin{array}{c} \frac{1}{2} + I_1, I_2 - \frac{n}{2}(q-1), I_3 - \frac{n}{2}(q-2), I_4 - \frac{n}{2}(q-3) \\ I_5 - \frac{n}{2}, I_6 + \frac{n}{2}, \frac{1}{2} + I_7 - \frac{n}{2}(q-2) \end{array} \middle| z \right) \ .
\end{aligned}
\qquad (125)
$$

Further simplification is possible for (i) $q = 3$ and (ii) $q = 4$. In these cases, we have:

- for $q = 3$,

$$
\begin{aligned}
& \{1, \theta\} \times {}_3F_2 \left(\begin{array}{c} 1, I_2 - \frac{n}{2}, \frac{1}{2} + I_3 + \frac{n}{2} \\ I_4 + n, \frac{1}{2} + I_5 \end{array} \middle| z \right) \ , \\
& \{1, \theta\} \times {}_2F_1 \left(\begin{array}{c} I_1 + \frac{1}{2} - \frac{n}{2}, I_2 - \frac{3}{2}n \\ I_3 + \frac{1}{2} - n \end{array} \middle| z \right) \ , \\
& \{1, \theta\} \times {}_3F_2 \left(\begin{array}{c} 1, \frac{1}{2} + I_1, I_2 - n \\ I_3 + \frac{n}{2}, I_4 + \frac{1}{2} - \frac{n}{2} \end{array} \middle| z \right) \ ,
\end{aligned}
\qquad (126)
$$

  and in accordance with Section 6.1, this integral has two master-integrals only

- for $q = 4$

$$
\begin{aligned}
& \{1, \theta, \theta^2\} \times {}_4F_3 \left(\begin{array}{c} 1, I_1 - \frac{n}{2}, I_2 - n, \frac{1}{2} + I_3 + \frac{n}{2} \\ I_4 + \frac{n}{2}, I_5 + n, \frac{1}{2} + I_6 - \frac{n}{2} \end{array} \middle| z \right) \ , \\
& \{1, \theta, \theta^2\} \times {}_3F_2 \left(\begin{array}{c} I_1 + \frac{1}{2} - \frac{n}{2}, I_2 - \frac{3}{2}n, I_3 - 2n \\ I_4 - \frac{n}{2}, I_5 + \frac{1}{2} + \frac{3}{2}n \end{array} \middle| z \right) \ , \\
& \{1, \theta, \theta^2\} \times {}_3F_2 \left(\begin{array}{c} \frac{1}{2} + I_1, I_2 - \frac{3}{2}n, I_3 - n \\ I_4 + \frac{n}{2}, I_5 + \frac{1}{2} - n \end{array} \middle| z \right) \ ,
\end{aligned}
\qquad (127)
$$

and in accordance with Section 6.1, this integral has three nontrivial master-integrals only (see [36]).

The three-loop diagram B^3_{1122} depicted in Fig. (2) corresponds to a particular case of diagram B^q_{11220} for $\sigma = 0$ and $q = 3$. In this case, $\Pi_{i=1}^{q-3} \frac{\Gamma(n/2-\sigma_i)}{\sigma_i} = 0$ and closing the contour of integration on the left we obtain:

$$
\begin{aligned}
&B^3_{1122}(m^2, M^2, \alpha_1, \alpha_2, \beta_1, \beta_2) = \frac{\left[i^{1-n}\pi^{\frac{n}{2}}\right]^3 (-m^2)^{\frac{n}{2}-\alpha_{1,2}}(-M^2)^{n-\beta_{1,2}}}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma(\beta_1)\Gamma(\beta_2)\Gamma\left(\frac{n}{2}\right)} \\
&\quad \left[\frac{\Gamma\left(\beta_1-\frac{n}{2}\right)\Gamma\left(\beta_2-\frac{n}{2}\right)\Gamma\left(\alpha_{1,2}-\frac{n}{2}\right)\Gamma\left(\frac{n}{2}\right)\Gamma(\alpha_1)\Gamma(\alpha_2)}{\Gamma(\alpha_{1,2})}\right. \\
&\quad \times {}_6F_5\left(\begin{array}{c} \alpha_1, \alpha_2, \alpha_{1,2}-\frac{n}{2}, \frac{n}{2}, \frac{1-\beta_{1,2}+n}{2}, \frac{2-\beta_{1,2}+n}{2} \\ 1-\beta_1+\frac{n}{2}, 1-\beta_2+\frac{n}{2}, 1-\beta_{1,2}+n, \frac{\alpha_{1,2}}{2}, \frac{\alpha_{1,2}+1}{2} \end{array}\middle| \frac{M^2}{m^2}\right) \\
&\quad + \frac{\Gamma\left(\frac{n}{2}-\beta_1\right)\Gamma\left(\frac{n}{2}-\beta_2\right)\Gamma\left(\beta_{1,2}-\frac{n}{2}\right)\Gamma\left(a_{\alpha_1,\beta_{1,2}}-n\right)\Gamma\left(a_{\alpha_2,\beta_{1,2}}-n\right)\Gamma\left(a_{\alpha_{1,2},\beta_{1,2}}-\frac{3n}{2}\right)}{\Gamma(\alpha_{1,2}+2\beta_{1,2}-2n)} \\
&\quad \times \left(\frac{M^2}{m^2}\right)^{\beta_{1,2}-n} {}_6F_5\left(\begin{array}{c} \beta_{1,2}-\frac{n}{2}, a_{\alpha_1,\beta_{1,2}}-n, a_{\alpha_2,\beta_{1,2}}-n, a_{\alpha_{1,2},\beta_{1,2}}-\frac{3n}{2}, \frac{1+\beta_{1,2}-n}{2}, \frac{2+\beta_{1,2}-n}{2} \\ 1+\beta_1-\frac{n}{2}, 1+\beta_2-\frac{n}{2}, 1+\beta_{1,2}-n, \frac{\alpha_{1,2}}{2}+\beta_{1,2}-n, \frac{\alpha_{1,2}+1}{2}+\beta_{1,2}-n \end{array}\middle| \frac{M^2}{m^2}\right) \\
&\quad + \frac{\Gamma(\beta_1)\Gamma\left(\beta_2-\frac{n}{2}\right)\Gamma\left(\frac{n}{2}-\beta_1\right)\Gamma\left(\alpha_1+\beta_1-\frac{n}{2}\right)\Gamma\left(\alpha_2+\beta_1-\frac{n}{2}\right)\Gamma\left(\alpha_{1,2}+\beta_1-n\right)}{\Gamma(\alpha_{1,2}+2\beta_1-n)} \\
&\quad \left(\frac{M^2}{m^2}\right)^{\beta_1-n/2} {}_6F_5\left(\begin{array}{c} \beta_1, \alpha_1+\beta_1-\frac{n}{2}, \alpha_2+\beta_1-\frac{n}{2}, \alpha_{1,2}+\beta_1-n, 1+\frac{\beta_1-\beta_2}{2}, \frac{1+\beta_1-\beta_2}{2} \\ 1+\beta_1-\beta_2, 1+\beta_1-\frac{n}{2}, 1-\beta_2+\frac{n}{2}, \frac{\alpha_{1,2}}{2}+\beta_1-n, \frac{\alpha_{1,2}+1}{2}+\beta_1-n \end{array}\middle| \frac{M^2}{m^2}\right) \\
&\quad + \frac{\Gamma(\beta_2)\Gamma\left(\beta_1-\frac{n}{2}\right)\Gamma\left(\frac{n}{2}-\beta_2\right)\Gamma\left(\alpha_1+\beta_2-\frac{n}{2}\right)\Gamma\left(\alpha_2+\beta_2-\frac{n}{2}\right)\Gamma\left(\alpha_{1,2}+\beta_2-n\right)}{\Gamma(\alpha_{1,2}+2\beta_2-n)} \\
&\quad \left.\left(\frac{M^2}{m^2}\right)^{\beta_2-n/2} {}_6F_5\left(\begin{array}{c} \beta_2, \alpha_1+\beta_2-\frac{n}{2}, \alpha_2+\beta_2-\frac{n}{2}, \alpha_{1,2}+\beta_2-n, \frac{1+\beta_2-\beta_1}{2}, \frac{2+\beta_2-\beta_1}{2} \\ 1+\beta_2-\beta_1, 1+\beta_2-\frac{n}{2}, 1-\beta_1+\frac{n}{2}, \frac{\alpha_{1,2}-n}{2}+\beta_2, \frac{\alpha_{1,2}+1-n}{2}+\beta_2 \end{array}\middle| \frac{M^2}{m^2}\right)\right].
\end{aligned} \tag{128}
$$

6.5 J^q_{023} & J^q_{012}

Let us consider the q-loop propagator type diagram J^q_{023} (see Fig. 3) defined as

$$
\begin{aligned}
&J^q_{023}(M_1^2, M_2^2, \alpha_1, \alpha_2, \sigma_1, \cdots, \sigma_{q-1}) \\
&= \int \frac{d^n(k_1 \cdots k_q)}{[k_1^2 - M_1^2]^{\alpha_1}[k_2^2 - M_2^2]^{\alpha_2}[k_3^2]^{\sigma_1} \cdots [(p - k_1 - \cdots - k_q)^2]^{\sigma_{q-1}}}\,.
\end{aligned} \tag{129}
$$

The Mellin-Barnes representation of Eq.(129) is

$$
\begin{aligned}
J^q_{023}(M_1^2, M_2^2, \alpha_1, \alpha_2, \sigma_1, \cdots, \sigma_{q-1}) = {} & \frac{[i^{1-n}\pi^{q/2}]^q}{\Gamma(\alpha_1)\Gamma(\alpha_2)} \left\{ \Pi_{i=1}^{q-1} \frac{\Gamma\left(\frac{n}{2}-\sigma_i\right)}{\Gamma(\sigma_i)} \right\} \\
(p^2)^{\frac{n}{2}q - a_{\alpha_1,\alpha_2,\sigma}} \int \int dt_1 dt_2 \Gamma(-t_1)\Gamma(-t_2) & \left(-\frac{M_1^2}{p^2} \right)^{t_1} \left(-\frac{M_2^2}{p^2} \right)^{t_2} \\
\times \frac{\Gamma\left(\frac{n}{2}-\alpha_1-t_1\right)\Gamma\left(\frac{n}{2}-\alpha_2-t_2\right)\Gamma\left(a_{\alpha_1,\alpha_2,\sigma}-n+t_1+t_2\right)}{\Gamma\left(\frac{3}{2}n - a_{\alpha_1,\alpha_2,\sigma}-t_1-t_2\right)} & \; , \qquad (130)
\end{aligned}
$$

where

$$
\sigma \;=\; \sum_{i=1}^{q-1} \sigma_i \; , \quad a_{\alpha_1,\alpha_2,\sigma} = \alpha_1+\alpha_2+\sigma \; , \quad a_{\alpha_2,\sigma} = \alpha_2+\sigma \; . \qquad (131)
$$

Closing the contour of integration in Eq.(130) on the left, we obtain

$$
\begin{aligned}
J^q_{023}(M_1^2, M_2^2, \alpha_1, \alpha_2, \sigma_1, \cdots, \sigma_{q-1}) = {} & \frac{[i^{1-n}\pi^{q/2}]^q (-M_1^2)^{\frac{n}{2}q - a_{\alpha_1,\alpha_2,\sigma}}}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma\left(\frac{n}{2}\right)} \left\{ \Pi_{i=1}^{q-1} \frac{\Gamma\left(\frac{n}{2}-\sigma_i\right)}{\Gamma(\sigma_i)} \right\} \\
\Bigg[\Gamma\left(\frac{n}{2}-\alpha_2\right) \Gamma\left(a_{\alpha_1,\alpha_2,\sigma}-\frac{n}{2}q\right) & \Gamma\left(a_{\alpha_2,\sigma}-\frac{n}{2}(q-1)\right) \\
\times F_4 \left(a_{\alpha_1,\alpha_2,\sigma} - \frac{n}{2}q, a_{\alpha_2,\sigma} - \frac{n}{2}(q-1), \frac{n}{2}, 1+\alpha_2-\frac{n}{2} \right| & \left. \frac{p^2}{M_1^2}, \frac{M_2^2}{M_1^2} \right) \\
+ \left(\frac{M_2^2}{M_1^2} \right)^{\frac{n}{2}-\alpha_2} \Gamma\left(\alpha_2 - \frac{n}{2}\right) \Gamma\left(a_{\alpha_1,\sigma}-\frac{n}{2}(q-1)\right) & \Gamma\left(\sigma-\frac{n}{2}(q-2)\right) \\
\times F_4 \left(a_{\alpha_1,\sigma} - \frac{n}{2}(q-1), \sigma - \frac{n}{2}(q-2), \frac{n}{2}, 1-\alpha_2+\frac{n}{2} \right| & \left. \frac{p^2}{M_1^2}, \frac{M_2^2}{M_1^2} \right) \Bigg] \; . \qquad (132)
\end{aligned}
$$

Let us analyze the result of the differential reduction of hypergeometric functions in Eq. (132) assuming that all powers of propagators are integer. For $q = 2$, both hypergeometric functions are reducible to function and their first derivative only (in accordance with the results of Section 3.6)

$$
\begin{aligned}
F_4(a+m_1, b+I_1, c+m_2, b|x,y) &= (1, \theta_x, \theta_y) \times F_4(a,b,c,b|x,y) + R_1(x,y) \; , \qquad (133) \\
F_4(a+m_1, I_1, c+m_2, b+m_3|x,y) &= (1, \theta_x, \theta_y) \times F_4(a,1,c,b|x,y) + R_2(x,y) \; , \qquad (134)
\end{aligned}
$$

where $\{m_i\}$ is a set of integers, and R_1, R_2 are rational functions. For $q \geq 3$ the result of differential reduction coincides with Eq. (77).

In accordance with Section 6.1, there are three nontrivial master integrals for $q = 2$ and four ones for $q \geq 3$.

For completeness, we also present the explicit results for the diagram J^q_{012} (see Fig. 3)(when external momenta on mass shell of one of the internal masses)

$$
\begin{aligned}
& J^q_{012}(m^2, M^2, \alpha, \lambda, \sigma_1, \cdots, \sigma_{q-1}) \\
& = \int \frac{d^n(k_1 \cdots k_q)}{[k_1^2 - m^2]^\alpha [k_2^2 - M^2]^\lambda [k_3^2]^{\sigma_1} \cdots [(p - k_1 - \cdots - k_q)^2]^{\sigma_{q-1}}} \Bigg|_{p^2=m^2} .
\end{aligned}
\tag{135}
$$

The hypergeometric representation for this diagram is

$$
\begin{aligned}
& J^q_{012}(m^2, M^2, \alpha, \lambda, \sigma_1, \cdots, \sigma_{q-1}) = \frac{[i^{1-n} \pi^{q/2}]^q (-M^2)^{\frac{n}{2}q - a_{\alpha,\lambda,\sigma}}}{\Gamma(\alpha)\Gamma(\lambda)\Gamma\left(\frac{n}{2}\right)} \left\{ \Pi_{i=1}^{q-1} \frac{\Gamma\left(\frac{n}{2} - \sigma_i\right)}{\Gamma(\sigma_i)} \right\} \\
& \Bigg[\Gamma\left(\frac{n}{2} - \alpha\right) \Gamma\left(a_{\alpha,\lambda,\sigma} - \frac{n}{2} q\right) \Gamma\left(a_{\alpha,\sigma} - \frac{n}{2}(q-1)\right) \\
& \quad \times {}_4F_3\left(\left. \begin{array}{c} a_{\alpha,\lambda,\sigma} - \frac{n}{2}q, a_{\alpha,\sigma} - \frac{n}{2}(q-1), \frac{\alpha}{2}, \frac{\alpha+1}{2} \\ 1 + \alpha - \frac{n}{2}, \alpha, \frac{n}{2}, \end{array} \right| \frac{4m^2}{M^2} \right) \\
& + \left(\frac{m^2}{M^2} \right)^{\frac{n}{2} - \alpha} \Gamma\left(\alpha - \frac{n}{2}\right) \Gamma\left(a_{\lambda,\sigma} - \frac{n}{2}(q-1)\right) \Gamma\left(\sigma - \frac{n}{2}(q-2)\right) \\
& \quad \times {}_4F_3\left(\left. \begin{array}{c} a_{\lambda,\sigma} - \frac{n}{2}(q-1), \sigma - \frac{n}{2}(q-2), \frac{n-\alpha}{2}, \frac{n-\alpha+1}{2} \\ \frac{n}{2}, n - \alpha, 1 - \alpha + \frac{n}{2} \end{array} \right| \frac{4m^2}{M^2} \right) \Bigg] ,
\end{aligned}
\tag{136}
$$

where

$$
\sigma \;=\; \sum_{i=1}^{q-1} \sigma_i \; , \quad a_{\alpha,\lambda,\sigma} = \alpha + \lambda + \sigma \; , \quad a_{\lambda,\sigma} = \lambda + \sigma \; .
\tag{137}
$$

In the case when the powers of propagators are all integer, the result of the differential reduction of the hypergeometric functions in Eq. (136) are the following (see also discussion in [37]):

- for $q = 2$

$$
\begin{aligned}
& \{1, \theta\} \times \; {}_2F_1\left(\left. \begin{array}{c} I_1 - n, \frac{1}{2} + I_2 \\ \frac{n}{2} + I_3 \end{array} \right| z \right) , \\
& \{1, \theta\} \times \; {}_3F_2\left(\left. \begin{array}{c} 1, I_1 - \frac{n}{2}, \frac{1}{2} + \frac{n}{2} + I_2 \\ n + I_3, \frac{n}{2} + I_4 \end{array} \right| z \right) ,
\end{aligned}
\tag{138}
$$

  In accordance with Eq.(6.1), there are two nontrivial master-integrals only.

- for $q \geq 3$

$$
\begin{aligned}
& \{1, \theta, \theta^2\} \times \; {}_3F_2\left(\left. \begin{array}{c} I_1 - \frac{n}{2}q, \frac{1}{2} + I_2, I_3 - \frac{n}{2}(q-1) \\ \frac{n}{2} + I_4, I_5 - \frac{n}{2} \end{array} \right| z \right) , \\
& \{1, \theta, \theta^2\} \times \; {}_3F_2\left(\left. \begin{array}{c} I_1 - \frac{n}{2}(q-2), I_2 - \frac{n}{2}(q-1), \frac{1}{2} + \frac{n}{2} + I_3 \\ n + I_4, \frac{n}{2} + I_5 \end{array} \right| z \right) ,
\end{aligned}
\tag{139}
$$

  In accordance with Eq.(6.1), there are three nontrivial master-integrals only.

7 Conclusion

The differential reduction algorithm [9] allows one to compare Horn-type hypergeometric functions with parameters whose values differ by integer. In this paper, we have presented the Mathematica based package **HYPERDIRE** for differential reduction of the generalized hypergeometric functions ${}_{p+1}F_p$ and Appell functions F_1, F_2, F_3 and F_4 to a set of basis functions, defined by Eqs. (19), (76), (77), respectively. These functions are closely related with a large class of Feynman Diagrams [5]. In contrast to existing packages, our package performs reduction of hypergeometric functions before ε-expansion and works with arbitrary values of parameters. The package could be easy extended to other Horn-type hypergeometric functions by adding new modules.

As an illustration of our approach, we considered a few examples, namely the Feynman diagrams shown in Figs. 1-3 with arbitrary powers of propagators and space-time dimension.

Acknowledgments.
We are grateful to A. Davydychev, A. Grozin, A. Isaev, A. Kotikov, O. Krotikova, G. Somogyi, V. Spiridonov, O. Tarasov, O. Veretin, B.F.L Ward, S. Yost for useful discussions, and G. Sandukovskaya for careful reading of the manuscript. M.Y.K.'s research was supported in part by BMBF, Grant No. 05H09GUE, DFG, Grant No. KN 365/3–2, and HGF, Grant No. HA 101.

References

[1] C.F. Gauss, Gesammelte Werke, vol. 3, Teubner, Leipzig, 1823, pp. 1866–1929.

[2] P. Appell and J. Kampé de Fériet, *Fonctions Hypergeometriques et Hyperspheriques. Polynomes d'Hermite* (Gauthier-Villars, Paris, 1926).

[3] W.N. Bailey, *Generalized Hypergeometric Series*, Cambridge University Press, 1935;
A. Erdelyi (Ed.), *Higher Transcendental Functions*, vol.1, McGraw-Hill, New York, 1953;
L.J. Slater, *Generalized hypergeometric functions*, Cambridge University Press, Cambridge 1966;
H. Exton, *Multiple hypergeometric functions and applications* John Wiley & Sons, Inc., New York-London-Sydney, 1976.
H.M. Srivastava, P.W. Karlsson, *Multiple Gaussian hypergeometric series*,
John Wiley & Sons, Inc., New York, 1985.

[4] I.M. Gel'fand, M.I. Graev, V.S. Retakh, Russian Math. Surveys **47** (1992) 1.

[5] C.G. Bollini,J.J. Giambiagi, Phys. Lett. B **40** (1972) 566; R. Gastmans, R. Meuldermans, Nucl. Phys. B **63** (1973) 277; D.S. Kershaw, Phys. Rev. D **8** (1973) 2708; A.C.T. Wu, Phys. Rev. D **9** (1974) 370; V.I. Kucheryavyi, Teor. Mat. Fiz. **20** (1974) 29; K. Mano, Phys. Rev. D **11** (1975) 452; R.J. Gonsalves, Phys. Rev. D

28 (1983) 1542; G. Kramer, B. Lampe, J. Math. Phys. **28** (1987) 945; E.E. Boos, A.I. Davydychev, Theor. Math. Phys. 89 (1991) 1052; A.I. Davydychev, J. Math. Phys. **32** (1991) 1052; A.I. Davydychev, J. Math. Phys. **33** (1992) 358; D.J. Broadhurst, Z. Phys. C **54**, 599 (1992); D.J. Broadhurst, J. Fleischer, O.V. Tarasov, Z. Phys. C **60** (1993) 287; A.I. Davydychev, J.B. Tausk, Nucl. Phys. B **397** (1993) 123; A.I. Davydychev, J.B. Tausk, Phys. Rev. D **53** (1996) 7381; F.A. Berends et al., Z. Phys. C **63** (1994) 227; S. Bauberger et al., Nucl. Phys. B **434** (1995) 383; F.A. Berends et all., Nucl. Phys. B **439** (1995) 536; A.V. Kotikov, Phys. Lett. B **375** (1996) 240; J. Fleischer et al., Nucl. Phys. B **539** (1999) 671 [Erratum-ibid. B **571** (2000) 511]; A.I. Davydychev, A.G. Grozin, Phys. Rev. D **59** (1999) 054023; T. Gehrmann, E. Remiddi, Nucl. Phys. B **580** (2000) 485; A.I. Davydychev, Phys. Rev. D **61** (2000) 087701; C. Anastasiou, E.W.N. Glover, C. Oleari, Nucl. Phys. B **575** (2000) 416 [Erratum-ibid. B **585** (2000) 763]; A.I. Davydychev, M.Yu. Kalmykov, Nucl. Phys. Proc. Suppl. **89** (2000) 283; O.V. Tarasov, Nucl. Phys. Proc. Suppl. **89** (2000) 237; L.G. Cabral-Rosetti, M.A. Sanchis-Lozano, J. Comput. Appl. Math. **115** (2000) 93 [arXiv:hep-ph/9809213]; A.I. Davydychev, M.Yu. Kalmykov, Nucl. Phys. B **605** (2001) 266; A.T. Suzuki, E.S. Santos, A.G.M. Schmidt, J. Phys. A **36** (2003) 4465; J. Fleischer, F. Jegerlehner, O.V. Tarasov, Nucl. Phys. B **672** (2003) 303; F. Jegerlehner, M.Yu. Kalmykov, O. Veretin, Nucl. Phys. B **658** (2003) 49; F. Jegerlehner, M.Yu. Kalmykov, Nucl. Phys. B **676** (2004) 365; T. Gehrmann, T. Huber, D. Maître, Phys. Lett. B **622** (2005) 295; A.I. Davydychev, Nucl. Instrum. Meth. A **559** (2006) 293; E. Bejdakic, Y. Schröder, Nucl. Phys. B (Proc. Suppl.) **160** (2006) 155; F. Jegerlehner, O.V. Tarasov, Phys. Lett. B **639** (2006) 299; O.V. Tarasov, Phys. Lett. B **638** (2006) 195; F. Jegerlehner, O.V. Tarasov, Nucl. Phys. Proc. Suppl. **157** (2006) 125; L.G. Cabral-Rosetti, M.A. Sanchis-Lozano, J. Phys. Conf. Ser. **37** (2006) 82 [arXiv:hep-ph/0206081]; A.G. Grozin, T. Huber, D. Maître; JHEP **0707** (2007) 033; J. Gluza, K. Kajda, T. Riemann, Comput. Phys. Commun. 177 (2007) 879; M.A. Shpot, J. Math. Phys. **48** (2007) 123512; O.V. Tarasov, Phys. Lett. B **670** (2008) 67; T. Huber, Nucl. Phys. B (Proc. Suppl.) 183 (2008) 238; V. Del Duca, C. Duhr, E.W. Nigel Glover, V.A. Smirnov, JHEP **1001**, 042 (2010); B.A. Kniehl, O.V. Tarasov, Nucl. Phys. B **820** (2009) 178; B.A. Kniehl, O.V. Tarasov, Nucl. Phys. B **833** (2010) 298; L.V. Bork, D.I. Kazakov, G.S. Vartanov, arXiv:1011.2440; G. Somogyi, arXiv:1101.3557;
T. Kaneko, arXiv: 1105.2080.

[6] E.D. Rainville, Bull. Amer. Math. Soc. **51** (1945) 714;
J. Wimp, Math. Comp. **22** (1968) 363;
W.Miller,Jr., SIAM J. Math. Anal., **3** (1972) 31; *ibid.* **4** (1973) 638; *ibid.* **5** (1974) 309;
J. Math. Phys. **13** (1972) 1393; SIAM J. Appl. Math., **25** (1973) 226;
Takeshi Sasaki, SIAM J. Math. Anal. **22** (1991) 821;
E. Horikawa, J. Math. Sci. Univ. Tokyo **1** (1994) 181;
A. Adolphson, B. Dwork, Trans. Amer. Math. Soc. **347** (1995) 615;
K. Roach, Proc. of ISSAC'96, 301-308, ACM, New York ;
K. Roach, Proc. of ISSAC'99, 205-211, ACM, New York;

R.G. Buschman, J. Comput. Appl. Math. **107** (1999) 127.

[7] J.A. Mullen, SIAM J. Appl. Math. **14** (1966) 1152.

[8] R.P. Singal, SIAM J. Math. Anal. **11** (1980) 390.

[9] N. Takayama, Japan J. Appl. Math. **6** (1989) 147.

[10] N. Takayama, J. Symbolic Comput., **20** (1995) 637.

[11] B. Sturmfels, N. Takayama, *Gröbner Bases and Hypergeometric Functions*, p. 246–258, London Math. Soc. Lecture Note Ser. 251, Cambridge Univ. Press, Cambridge, 1998;
M. Saito, B. Sturmfels, N. Takayama, Compositio Math. **115** (1999) 185–204;
M. Saito, B. Sturmfels, N. Takayama, *Gröbner Deformations of Hypergeometric Differential Equations*, (Springer-Verlag, Berlin, 2000).

[12] M. Bronstein, M. Petkovšek, Theoret. Comput. Sci. **157** (1996) 3–33;
F. Chyzak, B. Salvy, J. Symbolic Comput **26** (1998) 187-227;
F. Chyzak, *Gröbner Bases, Symbolic Summation and Symbolic Integration*, p.32–60, London Math. Soc. Lecture Note Ser. 251, Cambridge Univ. Press, Cambridge, 1998.

[13] M.Yu. Kalmykov, JHEP **0604** (2006) 056;
M.Yu. Kalmykov et al., Proc. of ACAT-08 [arXiv:0901.4716];
V.V. Bytev et al., [arXiv:0902.1352].

[14] V.V. Bytev, M.Yu. Kalmykov, B.A. Kniehl, Nucl. Phys. B **836** (2010) 129.

[15] M.Yu. Kalmykov, B.F.L. Ward, S. Yost, JHEP **0702** (2007) 040; JHEP **0711** (2007) 009;
S.A. Yost, M.Yu. Kalmykov, B.F.L. Ward, Proc. of ICHEP 2008 [arXiv:0808.2605];
M.Yu. Kalmykov et al., Proc. of Quarks-08, [arXiv:0810.3238] ;
M.Yu. Kalmykov, B. Kniehl, Nucl. Phys. B **809** (2009) 365; Phys. Part. Nucl. **41** (2010) 942.

[16] http://www.wolfram.com/

[17] S. Weinzierl, Comput. Phys. Commun. **145** (2002) 357;
S. Moch, P. Uwer, Comput. Phys. Commun. **174** (2006) 759.

[18] T. Huber, D. Maître, Comput. Phys. Commun. **175** (2006) 122; ibid **178** (2008) 755.

[19] J. Blumlein, S. Klein, C. Schneider F. Stan, arXiv:1011.2656.

[20] S. Moch, P. Uwer, S. Weinzierl, J. Math. Phys. **43** (2002) 3363;
S. Weinzierl, J. Math. Phys. **45** (2004) 2656.

[21] A.I. Davydychev, M.Yu. Kalmykov, Nucl. Phys. B **699** (2004) 3;
M.Yu. Kalmykov, Nucl. Phys. Proc. Suppl. **135** (2004) 280;
M.Yu. Kalmykov, B.F.L. Ward, S. Yost, JHEP **0710** (2007) 048.

[22] V.V. Bytev, M.Yu. Kalmykov, B.A. Kniehl, [arXiv:0904.0214v2], Chapter 4.

[23] M.Yu. Kalmykov, `http://theor.jinr.ru/~kalmykov/hypergeom/hyper.html`

[24] E. Bod, 'Algebraicity of the Appell-Lauricella and Horn hypergeometric functions", [arXiv:1005.0317].

[25] F. Beukers, "Monodromy of A-hypergeometric functions", [arXiv:1101.0493].

[26] P.W. Karlsson, J. Math. Phys. 12 (1971) 270.

[27] F.V. Tkachov, Phys. Lett. B 100 (1981) 65;
K.G. Chetyrkin, F.V. Tkachov, Nucl. Phys. B 192 (1981) 159.

[28] S. Laporta, Int. J. Mod. Phys. A **15** (2000) 5087;
C. Anastasiou, A. Lazopoulos, JHEP **0407** (2004) 046;
A.V. Smirnov, JHEP **0810** (2008) 107;
C. Studerus, Comput. Phys. Commun. **181** (2010) 1293.

[29] S.A. Yost et al., PoS **ICHEP2010** (2010) 135 [arXiv:1101.2348].

[30] O.V. Tarasov, Nucl. Phys. B **502** (1997) 455.

[31] A.A. Vladimirov, Theor. Math. Phys. 43 (1980) 417 [Teor. Mat. Fiz. 43 (1980) 210].

[32] G. 'tHooft and M. Veltman, Nucl. Phys. B **44** (1972) 189;
C.G. Bollini and J.J. Giambiagi, Nuovo Cim. B **12** (1972) 20;
J.F. Ashmore, Lett. Nuovo Cim. **4** (1972) 289;
G.M. Cicuta and E. Montaldi, Lett. Nuovo Cim. **4** (1972) 329.

[33] V.A. Smirnov, Commun. Math. Phys. **134** (1990) 109;
F.V. Tkachov, Int. J. Mod. Phys. A **8** (1993) 2047;
G.B. Pivovarov, F.V. Tkachov, Int. J. Mod. Phys. A **8** (1993) 2241.

[34] J. Fleischer M.Yu. Kalmykov, Comput. Phys. Commun. **128** (2000) 531;
F. Jegerlehner, M.Yu. Kalmykov, O. Veretin, Nucl. Phys. B **641** (2002) 285.

[35] L.V. Avdeev, Comput. Phys. Commun. **98** (1996) 15.

[36] S. Bekavac, A.G. Grozin, D. Seidel, V.A. Smirnov, Nucl. Phys. B **819** (2009) 183

[37] M.Yu. Kalmykov, B.A. Kniehl, in preparation.

[38] V.V. Bytev `https://sites.google.com/site/loopcalculations/home`